



Positive Quality Intervention: Selpercatinib Management

Description:

This PQI is developed to provide guidance for management of patients treated with selpercatinib (Retevmo).

Background:

Selpercatinib is a kinase inhibitor indicated for the treatment of:

- Adult patients with metastatic *RET* fusion-positive non-small cell lung cancer (NSCLC)
- Adult and pediatric patients 12 years of age and older with advanced or metastatic *RET*-mutant medullary thyroid cancer (MTC) who require systemic therapy
- Adult and pediatric patients 12 years of age and older with advanced or metastatic *RET* fusion-positive thyroid cancer who require systemic therapy and who are radioactive iodine-refractory (if radioactive iodine is appropriate)

This indication is approved under accelerated approval based on overall response rate and duration of response.¹

An efficacy study was conducted with patients with *RET* fusion-positive NSCLC who had received at least platinum-based chemotherapy; the objective response to selpercatinib of 64%. Median duration of response was 17.5 months and 63% of the responses were ongoing at a median follow-up of 12.1 months. For those who were previously untreated, ORR was measured at 85% (n=39).² Efficacy in patients with *RET*-mutant MTC who were previously treated with cabozantinib and/or vandetanib was studied and ORR was 69% (n=55) and median duration of response was not reached despite a median follow-up of 14.8 months. In this same study, cabozantinib or vandetanib-naïve patients had an ORR of 73% (n=88). *RET* fusion-positive thyroid cancer patients who were radioactive iodine (RAI)-refractory and were systemic therapy naïve had an ORR of 100% (CR 12.5%, PR 74%, n=8). Those thyroid patients previously treated with sorafenib, lenvatinib, or both had an ORR of 79% (CR 5.3%, PR 74%, n=19)³. Permanent discontinuation due to adverse reactions occurred in 5% of patients who have received selpercatinib. Dosage reductions due to an adverse reaction occurred in 31% of patients and the reactions where at least 2% or more of patients required reduction included ALT increased, AST increased, QT prolongation and fatigue.

PQI Process:

- Determine if a patient is eligible for selpercatinib
 - Review potential patients for presence of a *RET* gene fusion in NSCLC or thyroid cancer or presence of a *RET* gene mutation in MTC
 - Review the *Selpercatinib Genomic Testing Management PQI* for additional information

Upon prescription of selpercatinib

- Confirm Correct Dosing
 - Recommended dosage in adults and pediatric patients 12 years of age or older is based on weight
 - Less than 50 kg: 120 mg orally twice daily
 - 50 kg or greater: 160 mg orally twice daily
- Dosing considerations for Adverse Reactions
 - Recommended Selpercatinib Dose Reductions for Adverse Reactions¹

| Dose Reduction | Patients Weighing Less Than 50 kg | Patients Weighing 50 kg or Greater |
|----------------|-----------------------------------|------------------------------------|
| First | 80 mg orally twice daily | 120 mg orally twice daily |
| Second | 40 mg orally twice daily | 80 mg orally twice daily |
| Third | 40 mg orally once daily | 40 mg orally twice daily |

- Permanently discontinue if patient is unable to tolerate three dose reductions

Important notice: National Community Oncology Dispensing Association, Inc. (NCODA), has developed this Positive Quality Intervention platform. This platform represents a brief summary of medication uses and therapy options derived from information provided by the drug manufacturer and other resources. This platform is intended as an educational aid and does not provide individual medical advice and does not substitute for the advice of a qualified healthcare professional. This platform does not cover all existing information related to the possible uses, directions, doses, precautions, warning, interactions, adverse effects, or risks associated with the medication discussed in the platform and is not intended as a substitute for the advice of a qualified healthcare professional. The materials contained in this platform are for informational purposes only and do not constitute or imply endorsement, recommendation, or favoring of this medication by NCODA, which assumes no liability for and does not ensure the accuracy of the information presented. NCODA does not make any representations with respect to the medications whatsoever, and any and all decisions, with respect to such medications, are at the sole risk of the individual consuming the medication. All decisions related to taking this medication should be made with the guidance and under the direction of a qualified healthcare professional.



PQI Process Continued

- Recommended Selpercatinib Dose Reductions for Adverse Reactions¹

| Adverse Reaction | Severity | Dosage Modification |
|----------------------------|--------------------------|---|
| Hepatotoxicity | Grade 3 or Grade 4 | <ul style="list-style-type: none"> Withhold selpercatinib and monitor AST/ALT once weekly until resolution to Grade 1 or baseline. Resume at reduced dose by 2 dose levels and monitor AST and ALT once weekly until 4 weeks after reaching dose taken prior to the onset of Grade 3 or 4 increased AST or ALT. Increase dose by 1 dose level after a minimum of 2 weeks without recurrence and then increase to dose taken prior to the onset of Grade 3 or 4 increased AST or ALT after a minimum of 4 weeks without recurrence. |
| Hypertension | Grade 3 | <ul style="list-style-type: none"> Withhold selpercatinib for Grade 3 hypertension that persists despite optimal antihypertensive therapy. Resume at a reduced dose when hypertension is controlled. |
| | Grade 4 | <ul style="list-style-type: none"> Discontinue selpercatinib. |
| QT Interval Prolongation | Grade 3 | <ul style="list-style-type: none"> Withhold selpercatinib until recovery to baseline or Grade 0 or 1. Resume at a reduced dose. |
| | Grade 4 | <ul style="list-style-type: none"> Discontinue selpercatinib. |
| Hemorrhagic Events | Grade 3 or Grade 4 | <ul style="list-style-type: none"> Withhold selpercatinib until recovery to baseline or Grade 0 or 1. Discontinue selpercatinib for severe or life-threatening hemorrhagic events. |
| Hypersensitivity Reactions | All Grades | <ul style="list-style-type: none"> Withhold selpercatinib until resolution of the event. Initiate corticosteroids. Resume at a reduced dose by 3 dose levels while continuing corticosteroids. Increase dose by 1 dose level each week until the dose taken prior to the onset of hypersensitivity is reached, then taper corticosteroids. |
| Other Adverse Reactions | Grade 3 or Grade 4 | <ul style="list-style-type: none"> Withhold selpercatinib until recovery to baseline or Grade 0 or 1. Resume at a reduced dose. |

- Recommended Dosage for Concomitant Use of Strong and Moderate CYP3A Inhibitors

| Current Dosage | Recommended Dosage | |
|---------------------------|---------------------------|--------------------------|
| | Moderate CYP3A Inhibitor | Strong CYP3A Inhibitor |
| 120 mg orally twice daily | 80 mg orally twice daily | 40 mg orally twice daily |
| 160 mg orally twice daily | 120 mg orally twice daily | 80 mg orally twice daily |

- Recommended Dosage for Severe Hepatic Impairment

| Current Dosage | Recommended Dosage |
|---------------------------|--------------------------|
| 120 mg orally twice daily | 80 mg orally twice daily |
| 160 mg orally twice daily | 80 mg orally twice daily |

Important notice: National Community Oncology Dispensing Association, Inc. (NCODA), has developed this Positive Quality Intervention platform. This platform represents a brief summary of medication uses and therapy options derived from information provided by the drug manufacturer and other resources. This platform is intended as an educational aid and does not provide individual medical advice and does not substitute for the advice of a qualified healthcare professional. This platform does not cover all existing information related to the possible uses, directions, doses, precautions, warning, interactions, adverse effects, or risks associated with the medication discussed in the platform and is not intended as a substitute for the advice of a qualified healthcare professional. The materials contained in this platform are for informational purposes only and do not constitute or imply endorsement, recommendation, or favoring of this medication by NCODA, which assumes no liability for and does not ensure the accuracy of the information presented. NCODA does not make any representations with respect to the medications whatsoever, and any and all decisions, with respect to such medications, are at the sole risk of the individual consuming the medication. All decisions related to taking this medication should be made with the guidance and under the direction of a qualified healthcare professional.



Patient Centered Activities:

- Provide Oncology Chemotherapy Education (OCE) Sheet
- Counsel patient to swallow the capsules whole with or without food and to not crush or chew the capsules
- Review patient medication list to avoid concomitant use of strong and moderate CYP3A inhibitors
- Counsel patient to report adverse events related to high blood pressure, liver problems, heart rhythm changes, signs of bleeding, allergic reactions, and lack of wound healing
- Tumor lysis syndrome (TLS) occurred in 1% of patients with medullary thyroid carcinoma receiving selpercatinib
 - Counsel patient to contact care team to report signs and symptoms of TLS⁴
 - Nausea
 - Vomiting
 - Lack of appetite
 - Fatigue
 - Dark urine/flank pain
 - Reduced urine output
 - Numbness, seizures, hallucinations
 - Muscle cramps
 - Heart palpitations

References:

1. RETEVMO [package insert]. Lilly USA, LLC, Indianapolis, IN; Jan 2021
2. Drilon A, Oxnard GR, et al. Efficacy of Selpercatinib in *RET* Fusion–Positive Non–Small-Cell Lung Cancer. *N Engl J Med*. 2020; 383:813-824
3. Shah MH, Sherman EJ, et al. Selpercatinib (LOXO-292) in patients with *RET*-mutant medullary thyroid cancer. *J Clin Oncol*. 2020; 38 (15)
4. Gupta A, Moore JA. Tumor Lysis Syndrome. *JAMA Oncol*. 2018;4(6):895.

Important notice: National Community Oncology Dispensing Association, Inc. (NCODA), has developed this Positive Quality Intervention platform. This platform represents a brief summary of medication uses and therapy options derived from information provided by the drug manufacturer and other resources. This platform is intended as an educational aid and does not provide individual medical advice and does not substitute for the advice of a qualified healthcare professional. This platform does not cover all existing information related to the possible uses, directions, doses, precautions, warning, interactions, adverse effects, or risks associated with the medication discussed in the platform and is not intended as a substitute for the advice of a qualified healthcare professional. The materials contained in this platform are for informational purposes only and do not constitute or imply endorsement, recommendation, or favoring of this medication by NCODA, which assumes no liability for and does not ensure the accuracy of the information presented. NCODA does not make any representations with respect to the medications whatsoever, and any and all decisions, with respect to such medications, are at the sole risk of the individual consuming the medication. All decisions related to taking this medication should be made with the guidance and under the direction of a qualified healthcare professional.