

Darolutamide (Nubeqa®) in combination with Docetaxel (Taxotere) for Metastatic Castration-Sensitive Prostate Cancer

Description: This PQI aims to provide information on the administration, management of adverse events, patient follow-up, and recommended dose reductions for darolutamide in combination with docetaxel for metastatic hormone castration-sensitive prostate cancer (mCSPC).

Background:

Darolutamide is an androgen receptor inhibitor indicated for the treatment of adult patients with¹:

- mCSPC in combination with docetaxel and androgen deprivation therapy (ADT)
- mCSPC in combination with ADT
- non-metastatic castration-resistant prostate cancer (nmCRPC) in combination with ADT

Most common adverse reactions mCSPC in combination with docetaxel¹:

- >10% with a ≥2% increase over placebo: constipation, rash, decreased appetite, hemorrhage, increased weight, and hypertension
- Most common laboratory test abnormalities (≥30%): anemia, hyperglycemia, decreased lymphocyte count, decreased neutrophil count, increased AST, increased ALT, and hypocalcemia

PQI Process:

- Start ADT within 12 weeks before administering darolutamide and docetaxel
- Initiate darolutamide at 600 mg twice daily with food
 - For patients with severe renal impairment (eGFR 15-29 mL/min) the recommended dose is 300 mg BID
 - For patients with moderate hepatic impairment (Child-Pugh class B) recommended dose is 300 mg BID
- Start docetaxel IV 75mg/m² every 3 weeks for 6 cycles within 6 weeks of initiating darolutamide

Darolutamide dose management¹

- Treatment with darolutamide can be continued until disease progression/or unacceptable toxicity even if a dose of docetaxel is delayed, interrupted, or discontinued
- For severe renal impairment not receiving hemodialysis (GFR 15-29 mL/min/1.73 m²) reduce dose to 300 mg BID
- Hold darolutamide or dose reduce to 300 mg BID for grade 3 adverse reaction; may be resumed at 600 mg twice per day once the adverse reaction returns to baseline
- Doses under 300 mg twice a day are not recommended

Permanently discontinue darolutamide in the event of

- Grade 3-4 ischemic heart disease
- Development of seizures during darolutamide therapy

Clinical pearls

- Optimize cardiovascular risk factor management - hypertension, diabetes, dyslipidemia
- Use effective contraception during treatment and for 1week post last dose of darolutamide
- Avoid using darolutamide with a combined P-gp and strong or moderate CYP3A4 inhibitor/inducer

IMPORTANT NOTICE: NCODA has developed this Positive Quality Intervention platform. This platform is intended as an educational aid, does not provide individual medical advice, and does not substitute for the advice of a qualified healthcare professional. This platform does not cover all existing information related to the possible uses, directions, doses, precautions, warning, interactions, adverse effects, or risks associated with the medication. The materials contained in this platform do not constitute or imply endorsement, recommendation, or favoring of this medication by NCODA. NCODA does not ensure the accuracy of the information presented and assumes no liability relating to its accuracy. All decisions related to taking this medication should be made with the guidance and under the direction of a qualified healthcare professional. It is the individual's sole responsibility to seek guidance from a qualified healthcare professional. *Updated 7.25.25 PQI-134*

- If combination is necessary, monitor patient more frequently
- Review prescribing information of the BCRP, OATP1B3, OATP 1B1 substrates when used concomitantly with darolutamide

Docetaxel dose management⁴

- Administer drug when ANC is at least 1500 cells/mm³ or higher
- Dose reduce to 60 mg /m² if the patient experiences febrile neutropenia, ANC of <500 cells/mm³ for more than 1 week, severe or cumulative skin toxicities, or moderate neurotoxicity
- If the patient continues to experience the above side effects at 60 mg/m² the treatment should be discontinued
- Grade 3 liver dysfunction- reduce docetaxel by 20%

Permanently discontinue docetaxel in the event of:

- Grade 4 liver dysfunction
- Severe hypersensitivity reaction to docetaxel

Clinical pearls⁴⁻⁵

- Monitor for fluid retention, and manage per institution guidelines
- Be aware of the irritant/vesicant potential of docetaxel and consider central line in patients with poor IV access
- Pre-medicate with oral dexamethasone 8 mg twice daily x 3 days, starting the day before docetaxel administration
- Consider prophylactic pegfilgrastim 24 hours post docetaxel due to risk for febrile neutropenia⁶

Patient-Centered Activities:

- Provide darolutamide [Patient Education Sheet](#) and docetaxel [Patient Education Sheet](#)
- Side effects of combination therapy with darolutamide and docetaxel include neutropenia, neutropenic fevers, musculoskeletal pain, constipation, decreased appetite, rash, bleeding, weight gain, and hypertension
- Discuss risk for serious side effects including severe infusion reaction, development of seizures, and ischemic heart disease
- Ensure proper contraception and pregnancy protection is used and ensure the patient is aware that fertility may be impaired
- Provide education on temperature monitoring, whom to call if fever develops, and neutropenic precautions
- Recommend that patient immediately report any new or worsening symptoms
- Discuss the possible need for dose modifications of darolutamide, docetaxel, or both due to side effects
- Ensure the patient is aware of the use of steroids to prevent anaphylaxis and fluid retention associated with docetaxel

References:

1. Darolutamide prescribing information. Retrieved November 17, 2022.
2. Smith, M. H. (2022, (12)). Darolutamide and survival on metastatic hormone-sensitive prostate cancer. New England Journal of Medicine, pp. 1132-1142, DOI: 10.1056/nejmoa2119115.
3. National Comprehensive Cancer Network – Prostate Cancer. from https://www.nccn.org/professionals/physician_gls/pdf/prostate.pdf.
4. Docetaxel prescribing information. Retrieved November 17, 2022.
5. Jamal, E. Z. (2018). Primary prophylactic GCSF in patients receiving docetaxol-based chemotherapy for breast cancer. Journal of Clinical Oncology, 136:15.