

Fruquintinib (Fruzaqla®) for the management of colorectal cancer

Description: The purpose of this PQI is to review clinical information regarding fruquintinib (Fruzaqla®) treatment of metastatic colorectal cancer and to provide insights into best practices and treatment optimization

Background:

Fruquintinib is a selective, small-molecule kinase inhibitor of all three vascular endothelial growth factor (VEGF) receptors comprehensively inhibiting the VEGF pathway. Fruquintinib use is indicated in¹:

- Colorectal cancer
 - Metastatic, previously treated patients who have received fluoropyrimidine, oxaliplatin, and irinotecan-based chemotherapy, an anti-VEGF therapy, and, if RAS wildtype and medically appropriate, an anti-EGFR therapy

Most common adverse reactions (>20%)¹:

- Vascular: hypertension, hemorrhage
- Renal: proteinuria, increased serum creatinine
- Skin: palmar-plantar erythrodysesthesia (hand-foot syndrome)
- Respiratory: dysphonia
- Gastrointestinal: abdominal pain, diarrhea, stomatitis
- Miscellaneous: fatigue, musculoskeletal pain

PQI Process:

1. Eligibility Assessment and Clinical Criteria Confirmation:

- **Provider:** Confirm the diagnosis of mCRC and that the patient has previously received:
 - Fluoropyrimidine-, oxaliplatin- and irinotecan-based chemotherapy
 - Anti-VEGF therapy (i.e. bevacizumab in the first/second line setting)
 - If RAS wild-type and clinically indicated, anti-EGFR therapy
- Assess ECOG performance status to ensure appropriateness for fruquintinib initiation

2. Dosage Determination and Prescription Order:

- **Provider:** Determine the appropriate starting dose
 - 5 mg orally once daily for 21 days, followed by 7 days off
 - No dosage adjustment is recommended for patients with mild hepatic impairment; has not been sufficiently studied in patients with moderate hepatic impairment; not recommended for use in patients with severe hepatic impairment
- **Pharmacist:** Verify dose accuracy in the electronic health record and confirm consistency with clinical guidelines

3. Patient Education and Counseling:

- **Pharmacist/Provider:** Provide comprehensive education on fruquintinib, emphasizing:
 - The correct dosing regimen and adherence to the 21-day on, 7-day off cycle
 - Instructions on administration with or without food
 - The recognition and reporting of adverse events such as hypertension, hand-foot syndrome, and fatigue

IMPORTANT NOTICE: NCODA has developed this Positive Quality Intervention platform. This platform is intended as an educational aid, does not provide individual medical advice, and does not substitute for the advice of a qualified healthcare professional. This platform does not cover all existing information related to the possible uses, directions, doses, precautions, warning, interactions, adverse effects, or risks associated with the medication. The materials contained in this platform do not constitute or imply endorsement, recommendation, or favoring of this medication by NCODA. NCODA does not ensure the accuracy of the information presented and assumes no liability relating to its accuracy. All decisions related to taking this medication should be made with the guidance and under the direction of a qualified healthcare professional. It is the individual's sole responsibility to seek guidance from a qualified healthcare professional. *Updated 7.14.25 PQI-113*

- Reinforce the importance of follow-up visits and routine lab monitoring.
- 4. Drug Interaction Evaluation:**
 - **Pharmacist:** Conduct a complete medication reconciliation to identify potential interactions, particularly with CYP3A4 or inhibitors/inducers of CYP1A2, CYP2C19, CYP2C8, and CYP2C9
 - **Technician:** Assist with reviewing flagged medications for high-risk interactions and alert the pharmacist for further assessment
- 5. Prescription Review and Dispensing:**
 - **Provider:** Electronically prescribe fruquintinib with the correct regimen, including treatment duration and any necessary dose adjustments
 - **Pharmacist:** Validate the prescription against clinical standards and ensure that all necessary documentation is completed
 - **Technician:** Prepare, label, and package the medication accurately, adhering to pharmacy protocols
- 6. Lab Monitoring and Follow-Up Scheduling:**
 - **Provider:** Establish a follow-up plan for regular assessment of blood pressure, liver function tests (LFTs), proteinuria, and therapeutic response
 - **Pharmacist:** Schedule follow-up calls to monitor adherence and assess potential adverse effects.
 - **Technician:** Document and track lab results and escalate abnormal findings for review by the pharmacist and provider.
- 7. Adverse Event Management and Escalation:**
 - **Provider:** During follow-up visits, actively monitor patients for key adverse events such as hypertension, hand-foot syndrome, and LFT abnormalities
 - **Pharmacist:** Educate patients on identifying and reporting adverse events early and provide supportive care recommendations as needed. Recommend dose adjustments to provider as appropriate based on assessment of patient therapy tolerance including labs
 - **Technician:** Maintain accurate records of reported adverse events and communicate these to the pharmacist for further intervention

Patient-Centered Activities:

Provide Patient Education Materials:

Supply the patient with a [Patient Education Sheet](#) or equivalent educational material that explains fruquintinib's purpose, dosing regimen, potential side effects, and important safety information.

Counseling Pearls:

- **Administration Instructions:**
Instruct the patient to swallow tablets whole without crushing, chewing, or splitting them. If a dose is missed, it is ok to administer the dose if <12 hours have passed since scheduled dose but never take 2 doses on the same day to make up for missed dose.
- **If Vomiting Occurs:**
Advise the patient not to take an additional dose if vomiting occurs. They should continue with the next scheduled dose.
- **Hypertension Management:**
Educate the patient on regularly monitoring their blood pressure and reporting any increases to their healthcare team. Reinforce the importance of adhering to antihypertensive medications if prescribed.
- **Gastrointestinal Toxicity:**
Recommend taking fruquintinib with or without food as directed. For gastrointestinal symptoms

such as nausea or diarrhea, suggest the use of prophylactic or as needed antiemetics and over-the-counter antidiarrheal agents like loperamide, as needed.

- **Potential for Significant Drug Interactions:**

Encourage the patient to notify their healthcare team of any new medications, over-the-counter products, or herbal supplements they intend to use to avoid harmful drug interactions.

- **Hand-Foot Syndrome Precautions:**

Advise patients on preventive measures like using emollient creams, wearing comfortable shoes, and avoiding exposure to excessive heat or pressure on the palms and soles.

- **Photosensitivity Precautions:**

Recommend the use of sunscreen and protective clothing to minimize the risk of photosensitivity.

- **May Cause Reproductive Effects:**

Inform patients of child-bearing potential about the potential risks of infertility and the need for appropriate contraceptive measures during treatment.

References:

1. Lavacchi D, Roviello G, Guidolin A, et al. Evaluation of Fruquintinib in the Continuum of Care of Patients with Colorectal Cancer. *Int J Mol Sci.* 2023;24(6):5840. Published 2023 Mar 19. doi:10.3390/ijms24065840
2. Mumprecht V, Detmar M. Lymphangiogenesis and cancer metastasis. *J Cell Mol Med.* 2009;13(8A):1405-1416. doi:10.1111/j.1582-4934.2009.00834.x
3. Sun Q, Zhou J, Zhang Z, et al. Discovery of fruquintinib, a potent and highly selective small molecule inhibitor of VEGFR 1, 2, 3 tyrosine kinases for cancer therapy. *Cancer Biol Ther.* 2014;15(12):1635-1645. doi:10.4161/15384047.2014.964087
4. Fruzaqla (fruquintinib) [package insert]. Lexington, MA: Takeda Pharmaceuticals U.S.A., Inc.; 2023.
5. Li J, Qin S, Xu R, et al. Effect of fruquintinib vs placebo on overall survival in patients with previously treated metastatic colorectal cancer: the FRESCO randomized clinical trial. *JAMA.* 2018;319(24):2486-2496.
6. Dasari A, Lonardi S, Garcia-Carbonero R, et al; FRESCO-2 Study Investigators. Fruquintinib versus placebo in patients with refractory metastatic colorectal cancer (FRESCO-2): an international, multicenter, randomised, double-blind, phase 3 study. *Lancet.* 2023;402(10395):41-53. doi:10.1016/S0140-6736(23)00772-9

Supplemental Information:

Table 1. Recommended Dose Reductions for fruquintinib⁴

Dose Level	Fruquintinib dose
First dose reduction	4 mg orally once daily
Second dose reduction	3 mg orally once daily

Permanently discontinue fruquintinib in patients unable to tolerate 3 mg orally once daily