

# Isatuximab, Carfilzomib, and Dexamethasone (Isa-Kd)

Care Team Contact Information: \_\_\_\_\_

Pharmacy Contact Information: \_\_\_\_\_

Diagnosis: \_\_\_\_\_

- This treatment is often used for multiple myeloma, but it may also be used for other diagnoses.

Goal of Treatment: \_\_\_\_\_

- Treatment may continue for a certain time period, until it no longer works, or until side effects are no longer controlled.

## Treatment Regimen

- This treatment is often called by its acronym: Isa-Kd
  - **Isa:** Isatuximab
  - **K:** Carfilzomib (**K**yprolis)
  - **d:** Dexamethasone

Treatment Name	How the Treatment Works	How the Treatment is Given
Isatuximab (I-suh-TUK-sih-mab): Sarclisa (sar-KLIH-suh)	Helps your immune system find and attack cancer cells by targeting a specific protein on their surface	Infusion given into a vein.
Carfilzomib (kar-FIL-zoh-mib): Kyprolis (ky-PROH-lis)	Blocks a part of the cell that helps break down proteins, which stops cancer cells from growing and causes them to die.	Infusion given into a vein.
Dexamethasone (DEK-suh-MEH-thuh-son)	Tells cancer cells to "self-destruct".	Infusion given into a vein. OR Tablets taken by mouth.

**Treatment Administration and Schedule:** Treatment is typically repeated every 4 weeks. This length of time is called a “cycle”.

## Cycle 1:

- Dexamethasone is given on Days 1, 2, 8, 9, 15, 16, 22, and 23.
  - Note: Dexamethasone is typically given through a vein. However, it is often given by mouth on Day 23.
- Isatuximab is given weekly (Days 1, 8, 15, and 22).
- Carfilzomib is given on Days 1, 2, 8, 9, 15, and 16.
  - Note: You will receive a lower dose on Days 1 and 2. If tolerated, your care team will increase your dose.

Treatment Name	Cycle 1												
	Day 1	Day 2	...	Day 8	Day 9	...	Day 15	Day 16	...	Day 22	Day 23	...	Day 28
Dexamethasone	✓	✓		✓	✓		✓	✓		✓	✓		
Isatuximab	✓			✓			✓			✓			
Carfilzomib	✓	✓		✓	✓		✓	✓					

## Cycle 2 and Beyond:

- Dexamethasone is given on Days 1, 2, 8, 9, 15, 16, 22, and 23.
  - Note dexamethasone is typically given through a vein. However, it is often given by mouth on Days 22 and 23.
- Isatuximab is given every 2 weeks (Days 1 and 15).
- Carfilzomib is given on Days 1, 2, 8, 9, 15, and 16.

Treatment Name	Cycle 2													Cycle 3
	Day 1	Day 2	...	Day 8	Day 9	...	Day 15	Day 16	...	Day 22	Day 23	...	Day 28	Day 1
Dexamethasone	✓	✓		✓	✓		✓	✓		✓	✓			✓
Isatuximab	✓						✓							✓
Carfilzomib	✓	✓		✓	✓		✓	✓						✓

**Appointments:** Appointments may include regular check-ups with your care team, treatment appointments, lab visits, and imaging tests. It's important to keep your appointments whenever you can. If you miss any appointments, call your care provider as soon as possible to reschedule your appointment.

Supportive Care to Prevent and Treat Side Effects

Description	Supportive Care Given at the Clinic or Hospital	Supportive Care Taken at Home
To help prevent infusion reactions	<div></div> <div></div> <div></div>	<div></div> <div></div> <div></div>
To help prevent or treat nausea and vomiting	<div></div> <div></div> <div></div>	<div></div> <div></div> <div></div>
To help lower the risk of infections	<div></div> <div></div> <div></div>	<div></div> <div></div> <div></div>
Other	<div></div> <div></div> <div></div>	<div></div> <div></div> <div></div>

## Common Side Effects

Side Effect	Important Information
Low White Blood Cell (WBC) Count and Increased Risk of Infection	<p>Description: WBCs help protect the body against infections. If you have a low WBC count, you may be at a higher risk of infection.</p> <p>Recommendations:</p> <ul style="list-style-type: none"> <li>Wash your hands and bathe regularly.</li> <li>Avoid crowded places.</li> <li>Stay away from people who are sick.</li> <li>Your care team may prescribe a drug that promotes the growth of WBCs.</li> </ul> <p>Talk to your care team if you have:</p> <ul style="list-style-type: none"> <li>Fever of 100.4 °F (38°C) or higher</li> <li>Chills</li> <li>Cough</li> <li>Sore throat</li> <li>Painful urination</li> <li>Tiredness that is worse than normal</li> </ul>
Low Platelet Count	<p>Description: Platelets help the blood clot and heal wounds. If you have low platelet counts, you are at a higher risk of bruising and bleeding.</p> <p>Recommendations:</p> <ul style="list-style-type: none"> <li>Blow your nose gently and avoid picking it.</li> <li>Brush your teeth gently with a soft toothbrush and maintain good oral hygiene.</li> <li>Use an electric razor for shaving and a nail file instead of nail clippers.</li> <li>Avoid over-the-counter medications that may increase the risk of bleeding, such as NSAIDs.</li> <li>Talk with your care team or dentist before medical or dental procedures, as you may need to pause your treatment.</li> </ul> <p>Talk to your care team if you have:</p> <ul style="list-style-type: none"> <li>Nosebleed lasting over 5 minutes despite pressure</li> <li>Cut that continues to bleed</li> <li>Significant gum bleeding when flossing or brushing</li> <li>Severe headaches</li> <li>Blood in your urine or stool</li> <li>Blood in your spit after a cough</li> </ul>
Low Red Blood Cell (RBC) Count and Hemoglobin (Hgb)	<p>Description: RBCs and Hgb help bring oxygen to your body's tissues and take away carbon dioxide. If you have low RBC counts or Hgb, you may feel weak, tired, or look pale.</p> <p>Recommendations:</p> <ul style="list-style-type: none"> <li>Get 7 to 8 hours of sleep each night.</li> <li>Avoid operating heavy machinery when tired.</li> <li>Balance work and rest, staying active but resting when needed.</li> </ul> <p>Talk to your care team if you have:</p> <ul style="list-style-type: none"> <li>Shortness of breath</li> <li>Dizziness</li> <li>Fast or abnormal heartbeats</li> <li>Severe headache</li> </ul>
Trouble Breathing	<p>Description: Treatment may make you feel like it's a struggle to get enough air into your lungs.</p> <p>Recommendations:</p> <ul style="list-style-type: none"> <li>Stay hydrated by drinking plenty of fluids to help thin mucus.</li> <li>Avoid irritants such as smoke, strong odors, and allergens.</li> <li>Use humidifiers to add moisture to the air and soothe airways.</li> <li>Practice controlled breathing techniques, like pursed-lip breathing.</li> <li>Elevate your head with extra pillows when resting or sleeping, if needed.</li> <li>Limit physical exertion and take breaks during activities.</li> </ul> <p>Talk to your care team if you have:</p> <ul style="list-style-type: none"> <li>Shortness of breath</li> <li>Chest tightness</li> <li>Cough</li> </ul>

<p>High Blood Pressure</p>	<p>Description: Treatment can cause high blood pressure. It usually has no symptoms and can be dangerous if not treated. High blood pressure increases the risk of stroke, heart attack, and other health problems.</p> <p>Recommendations:</p> <ul style="list-style-type: none"> <li>• Exercise regularly, controlling your weight, and limiting alcohol and sodium intake.</li> <li>• If you are already being treated for high blood pressure, your care team may change your blood pressure medicine.</li> <li>• Your care team may ask you to track your blood pressure.</li> </ul> <p>Talk to your care team if you have:</p> <ul style="list-style-type: none"> <li>• Headaches</li> <li>• Dizziness or lightheadedness</li> <li>• Blurred vision</li> <li>• Trouble breathing</li> <li>• Nose bleeds</li> <li>• A pounding sensation in the chest, neck, or ears</li> <li>• Irregular heartbeats</li> <li>• Chest pain or pressure</li> </ul>
<p>Nausea or Vomiting</p>	<p>Description: Nausea is an uncomfortable feeling in your stomach or the need to throw up. This may or may not cause vomiting.</p> <p>Recommendations:</p> <ul style="list-style-type: none"> <li>• Eat smaller, more frequent meals.</li> <li>• Avoid fatty, fried, spicy, or highly sweet foods.</li> <li>• Eat bland foods at room temperature and drink clear liquids.</li> <li>• If you vomit, start with small amounts of water, broth, or other clear liquids when you are ready to eat again. If that stays down, then try soft foods (such as gelatin, plain cornstarch pudding, yogurt, strained soup, or strained cooked cereal). Slowly work up to eating solid food.</li> <li>• Your care provider may prescribe medicine for these symptoms.</li> </ul> <p>Talk to your care team if you have:</p> <ul style="list-style-type: none"> <li>• Vomiting for more than 24 hours</li> <li>• Vomiting that's nonstop</li> <li>• Signs of dehydration (like feeling very thirsty, having a dry mouth, feeling dizzy, or having dark urine)</li> <li>• Blood or coffee-ground-like appearance in your vomit</li> <li>• Bad stomach pain that doesn't go away after vomiting</li> </ul>
<p>Diarrhea</p>	<p>Description: Diarrhea is when you have loose, watery bowel movements more often than usual. The need to use the bathroom may occur urgently.</p> <p>Recommendations:</p> <ul style="list-style-type: none"> <li>• Keep track of how many times you go to the bathroom each day.</li> <li>• Drink 8 to 10 glasses of water or other fluids every day, unless your doctor tells you otherwise.</li> <li>• Eat small meals of mild, low-fiber foods like bananas, applesauce, potatoes, chicken, rice, and toast.</li> <li>• Stay away from foods with high fiber (like raw vegetables, fruits, and whole grains), foods that cause gas (like broccoli and beans), dairy foods (like yogurt and milk), and spicy, fried, and greasy foods.</li> <li>• Your care team might suggest a medicine for diarrhea.</li> </ul> <p>Talk to your care team if you have:</p> <ul style="list-style-type: none"> <li>• 4 or more bowel movements than normal in 24 hours</li> <li>• Dizziness or lightheadedness while having diarrhea</li> <li>• Bloody diarrhea</li> </ul>

Fatigue	<p>Description: Fatigue is a constant and sometimes strong feeling of tiredness.</p> <p>Recommendations:</p> <ul style="list-style-type: none"> <li>• Routine exercise has been shown to decrease levels of fatigue. Work with your care team to find the right type of exercise for you.</li> <li>• Ask your family and friends for help with daily tasks and emotional support.</li> <li>• Try healthy ways to feel better, like meditation, writing in a journal, doing yoga, and using guided imagery to lower anxiety and feel good.</li> <li>• Make a regular sleep schedule and limit naps during the day so you can sleep better at night, aiming for 7 to 8 hours of sleep.</li> <li>• Don't use heavy machines or do things that need your full attention if you're very tired to avoid accidents.</li> </ul> <p>Talk to your care team if you have:</p> <ul style="list-style-type: none"> <li>• Tiredness that affects your daily life</li> <li>• Tiredness all the time, and it doesn't get better with rest</li> <li>• Dizziness and weakness, along with being tired</li> </ul>
Infusion Reactions	<p>Description: Infusion reactions are common with isatuximab and can sometimes be severe or life threatening. They can also happen with carfilzomib but are less common.</p> <p>Recommendations:</p> <ul style="list-style-type: none"> <li>• Your care team will prescribe medicines before each infusion of isatuximab to help decrease your risk for infusion reactions or to help make any infusion reaction less severe.</li> <li>• You will be monitored for infusion reactions during each infusion of isatuximab.</li> <li>• Your care team may slow down or stop your infusion, or completely stop treatment with isatuximab if you have an infusion reaction.</li> </ul> <p>Get medical help right away if you develop any of the following symptoms of infusion reaction during or after an infusion:</p> <ul style="list-style-type: none"> <li>• Chills or shaking</li> <li>• Itching, rash, or flushing</li> <li>• Trouble breathing or wheezing; tongue-swelling</li> <li>• Dizziness or feeling faint</li> <li>• Fever of 100.4°F (or 38°C) or higher</li> <li>• Pain in your back or neck</li> </ul>

### Select Rare or Serious Side Effects

Side Effect	Talk to Your Care Team if You Have Any of These Signs or Symptoms		
Blood Clots	Signs or symptoms of a <b>blood clot</b> in the lung, arm, or leg may include: <ul style="list-style-type: none"> <li>• Shortness of breath</li> <li>• Chest pain</li> <li>• Arm or leg swelling</li> </ul>	Signs or symptoms of a <b>heart attack</b> may include: <ul style="list-style-type: none"> <li>• Chest pain that may spread to the arms, neck, jaw, back, or stomach area (abdomen)</li> <li>• Feeling sweaty</li> <li>• Shortness of breath</li> <li>• Feeling sick or vomiting</li> </ul>	Signs or symptoms of <b>stroke</b> may include: <ul style="list-style-type: none"> <li>• Sudden numbness or weakness, especially on one side of the body</li> <li>• Severe headache or confusion</li> <li>• Problems with vision, speech, or balance</li> </ul>
Heart Problems	<ul style="list-style-type: none"> <li>• Swelling of your stomach-area (abdomen), legs, hands, feet, or ankles</li> <li>• Shortness of breath</li> <li>• Nausea or vomiting</li> <li>• New or worsening chest discomfort, including pain or pressure</li> </ul>	<ul style="list-style-type: none"> <li>• Weight gain</li> <li>• Pain or discomfort in your arms, back, neck, or jaw</li> <li>• Protruding neck veins</li> <li>• Breaking out in a cold sweat</li> <li>• Feeling lightheaded or dizzy</li> </ul>	
Liver Problems	<ul style="list-style-type: none"> <li>• Yellowing of your skin or the whites of your eyes</li> <li>• Severe nausea or vomiting, pain on the right side of your stomach area (abdomen)</li> </ul>	<ul style="list-style-type: none"> <li>• Dark urine (tea colored)</li> <li>• Bleeding or bruising more easily than normal</li> </ul>	
Kidney Problems	<ul style="list-style-type: none"> <li>• Decrease in your amount of urine</li> <li>• Blood in your urine</li> </ul>	<ul style="list-style-type: none"> <li>• Swelling of your ankles</li> <li>• Loss of appetite</li> </ul>	
Brain Problems	<ul style="list-style-type: none"> <li>• Headache</li> <li>• Confusion or changes in the way you think,</li> </ul>	<ul style="list-style-type: none"> <li>• Seizures</li> <li>• Blurry vision or loss of vision</li> </ul>	
Herpes Reactivation	<ul style="list-style-type: none"> <li>• Blisters on your lips or around your mouth</li> <li>• Blisters on and around your genitals</li> </ul>	<ul style="list-style-type: none"> <li>• Fever of 100.4 °F (38°C) or higher</li> <li>• Flu-like symptoms, such as fatigue, headache, and muscle aches</li> </ul>	
Tumor Lysis Syndrome	<ul style="list-style-type: none"> <li>• Nausea or vomiting</li> <li>• Diarrhea</li> <li>• Muscle cramps or weakness</li> <li>• Tiredness</li> </ul>	<ul style="list-style-type: none"> <li>• Decreased amount of urine</li> <li>• Irregular heartbeat</li> <li>• Confusion</li> </ul>	

**Before starting treatment, ask your care team when to call 9-1-1 or seek emergency help.**  
**If you experience any new, worsening, or uncontrolled side effects, contact your care team immediately.**



## Intimacy, Pregnancy, and Breastfeeding

- Treatment may **change how you feel about intimacy and your body**. However, physical closeness—such as holding hands and hugging—remains safe. It is common to have questions about intimacy. If needed, talk to your care team for guidance.
- Treatment may **harm an unborn baby**.
  - If you are able to get pregnant, take a pregnancy test before starting treatment.
  - Use an effective method of birth control during treatment, for 5 months after your last dose of isatuximab, and for 6 months after your last dose of carfilzomib.
  - If you think you might be pregnant or if you become pregnant, tell your care team right away.
  - If your partner(s) could become pregnant, use an effective method of birth control—such as condoms—during treatment and for 3 months after your last dose of carfilzomib.
- **Do NOT breastfeed** during treatment and for 2 weeks after your last dose of carfilzomib.

## Handling Body Fluids and Waste

Some drugs you receive may stay in your urine, stool, sweat, or vomit for many days after treatment. Because many cancer drugs are toxic, your body waste may also be dangerous to touch. To help protect yourself, your loved ones, and the environment, **follow these instructions** for at least **48 hours** after each dose of **carfilzomib**:

- Pregnant women should avoid touching anything that may be soiled with body fluids from the patient.
- You can use your usual toilet. Always close the lid and flush to discard all waste. If you have a low-flow toilet, flush twice.
- If the toilet or seat is soiled with urine, stool, or vomit, clean the surface after each use before others use it.
- Wash your hands with soap and water for at least 20 seconds after using the toilet.
- If you need a bedpan, inform your caregiver so they can wear gloves and assist with cleanup. Wash the bedpan with soap and water daily.
- If you cannot control your bladder or bowels, use a disposable pad with a plastic back, a diaper, or a sheet to absorb waste.
- Wash any skin exposed to body waste with soap and water.
- Wash soiled linens or clothing separately from other laundry. If you don't have a washer, place them in a plastic bag until they can be washed.
- Wash your hands with soap and water after touching soiled linens or clothing.

## Additional Information

- **Tell your care team about all the medicines you take.**  
This includes prescriptions, over-the-counter drugs, vitamins, and herbal products. Before starting any new medicine, supplement, or vaccine, ask your care team first.
- **Treatment can affect the results of blood tests to match your blood type.**  
These changes can last for up to 6 months after your last dose of isatuximab. Your care team will do blood tests to match your blood type before you start treatment with isatuximab. Tell your care team that you are being treated with isatuximab before receiving blood transfusions.
- There is a risk of developing **new cancers** after taking isatuximab. Talk with your care team about the potential increased risk of new cancers.
- **This Patient Education Sheet may not describe all possible side effects.**  
Call your healthcare provider for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088.



## Notes

Updated Date: September 30, 2025

Scan the QR code below to access this education sheet.



Important notice: The Association of Cancer Care Centers (ACCC), Hematology/Oncology Pharmacy Association (HOPA), Network for Collaborative Oncology Development & Advancement, Inc. (NCODA), and Oncology Nursing Society (ONS) have collaborated in gathering information for and developing this patient education guide. This guide represents a brief summary of the medication derived from information provided by the drug manufacturer and other resources.

This guide does not cover all existing information related to the possible uses, directions, doses, precautions, warnings, interactions, adverse effects, or risks associated with this medication and should not substitute for the advice of a qualified healthcare professional. Provision of this guide is for informational purposes only and does not constitute or imply endorsement, recommendation, or favoring of this medication by ACCC, HOPA, NCODA, or ONS, who assume no liability for and cannot ensure the accuracy of the information presented. All decisions related to taking this medication should be made with the guidance and under the direction of a qualified healthcare professional.

Permission: Patient Education Sheets are provided as a free educational resource for patients with cancer and their caregivers in need of concise, easy to understand information about cancer therapy. Healthcare providers are permitted to copy and distribute the sheets to patients as well as direct patients to the Patient Education Sheets website. However, commercial reproduction or reuse, as well as rebranding or reposting of any type, are strictly prohibited without permission of the copyright holders. Permission requests, including direct linking from Electronic Health Records, and licensing inquiries should be emailed to [patienteducationssheets@ncoda.org](mailto:patienteducationssheets@ncoda.org).

Copyright © 2025 by Network for Collaborative Oncology Development & Advancement, Inc. All rights reserved.

PES-135