







THIOGUANINE

Name of your medication

Generic name — thioguanine (THY-oh-GWAH-neen) Brand name — Tabloid® (TAB-loyd)

Approved uses

Thioguanine is used to treat adult and pediatric patients with acute myeloid leukemia (AML).

Dose and schedule

Taking thioguanine as instructed is important to allow your treatment to be as effective as possible, so here are some key points to remember.

- ☐ Your dose is based on many factors, including your height and weight, overall health, and diagnosis.
- ☐ Thioguanine can be taken with or without food, but at the same time each day.
- Thioguanine should be taken whole and not crushed, cut, or dissolved. If you are unable to swallow thioguanine, talk to your care provider or pharmacist for possible options.
- If you miss a dose of thioguanine, **do not** take an extra dose or two doses at one time. Simply take your next dose at the regularly scheduled time. Be sure to write down if you miss a dose, and let your care provider know about any missed doses.

Drug and food interactions

- ☐ Thioguanine has many drug interactions, please inform your care provider of all prescription medications, over-the counter medications, vitamins, and herbal products that you are taking.
- ☐ Talk with your care provider or pharmacist before taking new medications or supplements, or receiving any vaccines.

Storage and handling

Handle thioguanine with care. Just like when chemotherapy is given into the vein, this drug can be toxic, and exposure of the drug to others should be limited.

- □ Store thioguanine at room temperature (68°F–77°F) in a dry location away from light.
- Keep thioguanine out of reach of children and pets.
- ☐ Leave thioguanine in the provided packaging until it is ready to be taken.
- Whenever possible, you should give thioguanine to yourself and follow the steps below. If a family member, friend, or caregiver needs to give the thioguanine to you, they also need to follow these steps.
 - 1. Wash hands with soap and water.
 - 2. Put on gloves to avoid touching the medication. (Gloves are not necessary if you give the drug to yourself.)
 - 3. Gently transfer the thioguanine from its package to a small medicine or other disposable cup.
 - 4. Administer the medicine immediately by mouth with water.
 - 5. Remove gloves and do not use them for anything else.









Continued on the next page

THIOGUANINE

- 6. Throw gloves and medicine cup in household trash.
- 7. Wash hands with soap and water.
- ☐ If a daily pill box or pill reminder will be used, contact your care team before using:
 - The person filling the box or reminder should wear gloves (gloves are not necessary if you are filling the box or reminder).
 - When empty, the box or reminder should be washed with soap and water before refilling. Be sure to wash hands with soap and water after the task is complete, whether or not gloves are worn
- If you have any unused thioguanine, **do not** throw it in the trash and **do not** flush it down the sink or toilet. Talk to your care provider or pharmacist about proper disposal of thioguanine.
- If you are traveling, put your thioguanine's packaging in a sealed plastic bag. Ask your pharmacist if any additional travel precautions are needed.

Side Effects of Thioguanine

Below are common side effects that have been known to happen in about one third or more of patients taking thioguanine are listed on the left side of this table. You **MAY NOT** experience these side effects. Options to help manage any side effects that do occur are included on the right side of this table. These should be discussed with your care provider. If you experience any side effect you cannot manage or that is not listed here, contact your care provider.

Possible Side Effect	Management
Decreased white blood cells (WBCs) and increased risk for infection	Your WBCs should be monitored by a simple blood test. When your WBCs are low, you are at a greater risk of having an infection. Take the following precautions to protect yourself from infection. • Wash your hands often, especially before eating and after using the bathroom. • Avoid crowds and people with fevers, flu, or other infection. • Bathe regularly to keep good personal hygiene. Contact your care provider if you experience any signs or symptoms of an infection such as: • Fever (temperature more than 100.4°F or 38°C) • Chills • Sore throat • Burning with urination • Unusual tiredness • A sore that becomes red, is draining, or does not heal Check with your care provider before taking any medicine for a fever or chills.









THIOGUANINE

Possible Side Effect	ssible Side Effect Management	
Decreased platelet count and increased risk of bleeding	Your platelets should be monitored by a simple blood test. When they are low, you may bruise or bleed more easily than usual. Use caution to avoid bruises, cuts, or burns. Blow your nose gently, and do not pick your nose. Brush your teeth gently with a soft toothbrush, and maintain good oral hygiene. When shaving, use an electronic razor instead of razor blades. Use a nail file instead of nail clippers. Call your care provider if you have bleeding that won't stop. Examples include: A bloody nose that bleeds for more than 5 minutes despite pressure A cut that continues to ooze despite pressure Gums that bleed excessively when you floss or brush Seek medical help immediately if you experience any severe headaches, observe blood in your urine or stool, cough up blood, or experience prolonged and uncontrollable bleeding. You may need to take a break or hold your medication for medical or dental procedures. Talk to your care provider or dentist before any scheduled procedures.	
Decreased hemoglobin, part of the red blood cells that carry iron and oxygen	Your hemoglobin should be monitored by a simple blood test. When your hemoglobin is low, you may notice that you get tired or fatigued more easily. Try to get 7–8 hours of sleep per night. Avoid operating heavy machinery if you feel too tired. Find a balance between work and rest. Stay as active as possible, but know that it is okay to rest as needed. You might notice that you are more pale than usual. Let your care provider know right away if you experience: Shortness of breath Dizziness Palpitations	
Changes in liver function	Your liver function should be checked periodically by a simple blood test. Contact your care provider if you notice any of the following: • Yellowing of the skin or whites of your eyes • Dark or brown urine • Unusual bleeding or bruising	









THIOGUANINE

Serious side effects of thioguanine

You may have a mild to moderate rise in blood pressure while taking thioguanine	Take your blood pressure on a regular
basis and let your doctor know of any big increases in blood pressure.	

- ☐ Thioguanine may be harmful to your liver. Speak to your care provider to know when you need to have laboratory tests done to monitor your liver function. If your get this side effect, your doctor may change your dose or stop treatment for some time.
- ☐ While taking thioguanine your skin may be more sensitive to the sun. Avoid being out in the sun when possible and cover your skin if you are out in the sun.

If you experience ANY uncontrolled side effect, call your physician or healthcare center immediately:

_____ (INSTITUTIONAL CONTACT INFO)

Handling body fluids and waste

Since thioguanine remains in your body for several days after it is taken, some of the drug may be present in urine, stool, sweat, or vomit. Once you have started to take thioguanine, it is important to know the following instructions every day for as long as your treatment lasts. This is to keep yourself, loved ones, and the environment as safe as possible.

- ☐ Pregnant women should avoid touching anything that may be soiled with body fluids from the patient.
- □ Toilet and septic systems
 - You may use the same toilet, septic tank, and/or sewer that you usually use. If you have a low-flow toilet, close the lid and flush twice to ensure all waste has been discarded.
 - If the toilet or toilet seat becomes soiled with urine, stool, or vomit, clean the surfaces before other people use the toilet.
 - Wash hands with soap and water after using the toilet.
- ☐ If you need a bedpan, be sure your caregiver knows to wear gloves to assist with cleanup and to wash the bedpan with soap and water every day.
- ☐ If you do not have good control of bladder or bowels, use a disposable pad with a plastic back, a diaper, or a sheet to absorb body waste.
- ☐ Wash any skin that has been exposed to body waste or thioguanine with soap and water.
- Linens or clothing that are soiled with body fluids or body waste should be washed separately from other linens and clothing. If you do not have a washer, place the soiled linens in a plastic bag until they can be washed.
- ☐ Wash hands with soap and water after touching linens or clothing that may be soiled with body fluids.









THIOGUANINE

Jraananav aaviia	I AATIVIITVI ABA	LAANTKAAANI	11 A 10
Pregnancy, sexual	i aciiviiv and	i COntraceoi	

Pregr	nancy, sexual activity, and contraception
	Women should not become pregnant and men should not get a partner pregnant while taking thioguanine. Men and women of childbearing age and potential should use effective contraception during therapy and for a minimum of six months after the last dose of thioguanine.
	Effective contraception could include one or more of the following: oral contraceptive, barrier methods, etc.
	Do not breastfeed while taking thioguanine and for one month after the last dose of thioguanine.
	Please inform your care provider if you become pregnant.
	It is safe to hug and kiss. Special precautions may be needed for sexual activity while on oral chemotherapy, and you are encouraged to ask your care provider.
Obtai	ning medication
Talk	with your care provider about the process for obtaining your thioguanine.
	(PHARMACY OR SPECIALTY PHARMACY CONTACT INFO)
Addit	ional resources
Pr	oduct prescribing information: www.accessdata.fda.gov/drugsatfda_docs/label/2004/12429s022lbl.pdf
	Updated — March 10, 2020
Addit	ional instructions

Important notice: The Association of Community Cancer Centers (ACCC), Hematology/Oncology Pharmacy Association (HOPA), National Community Oncology Dispensing Association, Inc. (NCODA), and Oncology Nursing Society (ONS) have collaborated in gathering information for and developing this patient education guide. This guide represents a brief summary of the medication derived from information provided by the drug manufacturer and other resources. This guide does not cover all existing information related to the possible uses, directions, doses, precautions, warnings, interactions, adverse effects, or risks associated with this medication and should not substitute for the advice of a qualified healthcare professional. Provision of this guide is for informational purposes only and does not constitute or imply endorsement, recommendation, or favoring of this medication by ACCC, HOPA, NCODA, or ONS, who assume no liability for and cannot ensure the accuracy of the information presented. The collaborators are not making any representations with respect to the medications whatsoever, and any and all decisions, with respect to such medications, are at the sole risk of the individual consuming the medication. All decisions related to taking this medication should be made with the guidance and under the direction of a qualified healthcare professional.

Permission: Oral Chemotherapy Education (OCE) sheets are provided as a free educational resource for patients with cancer in need of concise, easy-to-understand information about oral cancer drugs. Healthcare providers are permitted to copy and distribute the sheets to patients as well as direct patients to the OCE website for information. However, commercial reproduction or reuse, as well as rebranding or reposting of any type, are strictly prohibited without permission of the copyright holder. Please email permission requests and licensing inquiries to Contact@NCODA.org.

Copyright © 2019 by Hematology/Oncology Pharmacy Association. All rights reserved.