




1




2




3

[illegible]


2



7



8



9



Perceived CAR-T Barriers Among US-based Large Academic and Community Practice

Barriers		Potential Solutions	
Reimbursement/ Financial burden	- Health plan restrictions/denials for coverage of adoptive cell therapies - Slow approval process by payers - Patient deterioration (ineligible)	Reimbursement/ Financial burden	- Streamline mechanisms for equitable reimbursement - Utilize manufacturer-provided resources for patient support, insurance coverage and benefit verification

Aschall R, et al. Transplant Cell Ther. 2024;30(2S): S3703286. Gupta A, et al. Immunotherapy. 2020 Jul 12(10): 725-732.

Championing Medically Integrated Oncology: CELEBRATING A DECADE OF IMPACT

10



Advocacy for Safe and Equitable Access: Mission Accomplished June 26, 2025

- ASTCT 80/20 Taskforce (now subcommittee) was developed in 2020
 - Standardize/streamline requirements for onboarding cellular therapies
- Mission: provide safe use of CAR-T
 - Collaborative burden reduction

FDA NEWS RELEASE

FDA Eliminates Risk Evaluation and Mitigation Strategies (REMS) for Autologous Chimeric Antigen Receptor CAR T cell Immunotherapies

Agency determines the safety and effectiveness of these immunotherapies can be assured without a REMS

ASTCT 80/20 effectively advocated to streamline

ASTCT 80/20 consensus on key cell therapy community efforts to safe delivery of CAR-T

responsibility for standards of care in CAR-T therapies

provide CAR-T safety education

used to a central repository, such as each to each manufacturer

al accreditation groups, such as FACT, that oversight, not manufacturers

and experience have different needs for oversight

Lecker F, et al. Transplant Cellular Ther. 2025;31:345.e1-345.e2. Image available under creative commons CC BY-NC-ND license. FDA Eliminates Risk Evaluation and Mitigation Strategies (REMS) for Autologous Chimeric Antigen Receptor CAR T cell Immunotherapies. Accessed August 18, 2025, from: <https://www.fda.gov/news-events/press-announcements/fda-eliminates-risk-evaluation-and-mitigation-strategies-terms-autologous-chimeric-antigen-receptor>

Championing Medically Integrated Oncology: CELEBRATING A DECADE OF IMPACT

11



Financial Considerations for OP CAR-T: Creating a Sustainable and Patient-centric Approach

Reimbursement varies based upon public vs. private insurance and administration location

- High cost of CAR-T can be exacerbated by reimbursement restrictions
 - Ensure single case/contractual agreements include IP and OP sites of care
 - OP reimbursement*: average \$ for CAR-T + ~ 6%
 - Medicare uses OP prospective payor system (OPPS):
 - PPS-non-exempt: 99% of other centers (fixed rate per patient) vs. PPS-exempt: 11 Cancer Centers ("reasonable cost payment")
 - IP reimbursement MS-DRG base payment
 - Medicare copays capped
 - Medicaid coverage varies (product/state-specific)

Ensure thorough EMR documents for labeling criteria

*Product price increase may occur and need to be vigilantly monitored for updates
MS-DRG Medicare Severity Diagnostic-Related Group

Aggarwalli S, et al. Oncol Ther. 2023 Jun 1(2):283-276. American Society for Transplantation and Cellular Therapy. ASTCT CAR-T therapy coding and billing guide. Updated January 2025. Accessed August 18, 2025, <https://www.astct.org/docs/default-source/20250601-car-t-therapy-coding-and-billing-guide.pdf>. Hines M, et al. Future Oncol. 2023;21(10):1750-1759.

Championing Medically Integrated Oncology: CELEBRATING A DECADE OF IMPACT

12



Financial Considerations for OP CAR-T: Creating a Sustainable and Patient-centric Approach

Even with insurance there is significant financial burden

- Medical costs, lost wages (patient/caregiver) and lodging/transportation
- Ensure access to general financial support for these out-of-pocket expenses
 - Local lodging programs or grants

Shifting CAR-T to the OP setting may improve patient experience, optimize HCRU, and decrease costs:

- Decreased hospital bed usage
- Shorter hospital stays
- Lower costs (patient and institution)
- Improved patients QoL

Jagannath S, et al. *Clinical Ther*. 2023 Jun;11(2):269-275. American Society for Transplantation and Cellular Therapy. ASTCT CAR-T Therapy: setting and delivery issues. *Transpl January 2020*. Accessed August 19, 2025. <https://doi.org/10.1016/j.jct.2020.01.001>. Alanna M, et al. *Future Oncol*. 2023;19(11):1131-1138.

HCRU: Healthcare resource utilization; QoL: quality of life; EMR: electronic medical record


Championing Medically Integrated Oncology: CELEBRATING A DECADE OF IMPACT

13



OP Administration Can Reduce Healthcare and Patient Costs While Maintaining Suitable Benefit-Risk Profile


Jagannath S, et al. Citta-cel Component Cost of CAR-T: Average Total Costs Per Patient Based on 3 Phases*



Phase	100% IP	85% IP/15% OP	70% IP/30% OP
Pre-infusion	\$160,933	\$158,095	\$155,257
Peri-infusion	\$23,154	\$20,316	\$17,478
Post-infusion	\$18,268	\$4,584	\$4,584
Total	\$180,933	\$158,095	\$155,257

*Population based on CARTITUDE-1: Exploring 3 Phases of Cost Components: pre-infusion, peri-infusion, post-infusion


Palomba ML, et al. Liso-cel Cost of Care: IP vs. OP



Retrospective review of TRANSCEND 001 and OUTREACH trials (N = 303)

- 62% infused OP required admission
 - 21% within 72 hours
 - Admission LOS, median (range): 12 (0-88) IP vs. 4 (0-77) OP
- 38% of OP never required admit within 6-months
- Estimated 6-month post-CAR-T HCRU savings: \$89,535 IP vs. \$36,702 OP

Hansen, et al. Direct Costs of pooled TRANSCEND 001, OUTREACH, TRANSFORM and PILOT (liso-cel) Clinical Trials with OP and IP data



6-month post-infusion cost 2.4x higher
Cost reduction driven by reduced hospitalization costs

14



Considerations for Successful Implementation of Outpatient CAR-T



2024 NCODA INTERNATIONAL FALL SUMMIT

15

QUESTION ARS #1

2025 NCODA INTERNATIONAL FALL SUMMIT

You are currently planning on developing outpatient CAR-T at your institution. During a strategic planning meeting you were asked what components of CAR-T can be administered in outpatient care setting?

a. Lymphodepleting chemotherapy (Day -5 to Day -3)

b. Cells infusion (D0)

c. Close monitoring post-cell infusion for certain eligible products

d. All of the above

Championing Medically Integrated Oncology: CELEBRATING A DECADE OF IMPACT

16

Suitability Criteria for OP Administration of CAR-T

Key: Purposeful Selection



Patient

- Limited comorbidities with good PS

- Reliable 24-hour caregiver support

- Relatively stable disease burden, low risk of severe CRS/CANS



Product

- Predictable timeline of toxicities

- Gradual onset of potentially severe toxicities

Myers GD, et al. J Immunother Cancer. 2021;9:e402055. | Oliveira OC, et al. Transplant Cellular Ther. 2024, Feb;30(2):131-142.

PS: performance status



17

Overall Patient/Product Considerations for CAR-T Eligibility*

Indications and limitations

How time sensitive/urgent is treatment?

Is the patient at high risk of toxicity?

Consider antigen loss (e.g., CD19 for CD19-directed CAR-T)

Insurance coverage/preference of product

High tumor burden/need for bridging therapy

Multiple comorbidities

Life expectancy > 6 weeks

Adequate organ function

ECOG PS: 0-2

Ability to tolerate LD chemo

Ability to tolerate CAR-T cell-related toxicities

No active/ uncontrolled infections

Caregiver support pre, during, and post CAR-T

Housing/financial support

*Criteria for eligibility may differ from site by site

ECOG PS: Eastern Cooperative Oncology Group performance score

Alvarado JS, et al. Am Soc Clin Oncol Educ Book. 2019;38:448-453. | Yee BC, et al. Hematology. 2019;10(2):287-316.

Championing Medically Integrated Oncology: CELEBRATING A DECADE OF IMPACT

18

NCODA: Fall Summit

6

19

20

21

Systemic Spectrum of CAR-T Associated Toxicities

ICANS

- Confusion
- Reduced responsiveness
- Seizures
- General edema

TIAN

- Cerebral edema
- Increased ICP*
- Localized neural dysfunction

Cocircopathy

- Bleeding
- Thromboses

ICAH

- Anemia
- Thrombocytopenia
- Neutropenia

CRS

- Fever
- Hypertension
- Hypoxia

IEC-HS

- Hypertension
- Elevated LFTs
- Hemophagocytosis
- Cytopenias
- Cocircopathy

Infections

- Viral, bacterial, fungal
- Increased risk with cytopenias, B-cell depletion, and anti-CRS management

Clinical Symptoms/Findings

Neurologic/Muscular/Skeletal

- Headache
- Non-ICANS neurotoxicity
- Myalgia/Arthralgias

Pulmonary

- Dyspnea
- Tachypnea
- Hypoxia

General (flu-like)

- Fatigue
- Nausea/Vomiting
- Fevers/Rigors

TIAN: tumor inflammatory-associated neurotoxicity; ICP: intracranial pressure; IEC-HS: Immune effector cell associated hemophagocytic lymphohistiocytosis-like syndrome; LFT: liver function test; ICAH: Immune Effector Cell-Associated Hematotoxicity

Reis, A.W. et al. Cancer Medicine 2023; 12: 17. Image available at <https://www.researchsquare.com/publication/1234567890>. Reis, A.W. et al. Transplant Cell Ther 2023; 29(4):438-449. Jan, Y. et al. Blood 2023; 141(25):2482-2493. Cohen, D.S. et al. Lancet Oncol 2020; 21: 4020-4033. Saito, M. et al. J Immunother Cancer 2022; 10(12):e0430111. Gershwin, R.B. et al. J Clin Oncol 2021; 39:3620-3630. 2020; 38(25):2482-2493. Cohen, D.S. et al. Lancet Oncol 2020; 21: 4020-4033. Lee, D.H. et al. Biol Blood Marrow Transplant 2019; 25:1029.

22

CAR Construct Directly Associated with Onset/Severity of AE's

Domain Role Call:

- Antigen Binding Domain: Recognizes CD19 or BCMA antigen on B-cells
- Costimulatory Domain: Increase T-cell activation and enhances cytolytic function of T-cells
- CD3-zeta chain signaling domain: Induces T-cell activation

Diagram illustrating the CAR construct domains: 1. CD-19 or BCMA Antigen Binding, 2. Costimulatory Domain (4-1BB, CD28), 3. Signaling Domain (CD3ζ, CD28).

CD28	4-1BB
Member of the immunoglobulin (Ig) family	Member of the TNF receptor superfamily
Expands preferentially effector-like T-cells rapidly (limited CAR persistence)	Expands memory-like T-cells (slow activation but enhanced persistence)

SCFV: short-chain variable fragment, TM: transmembrane

Line graph showing CAR T cell count over 12 months for CD28 and 4-1BB constructs. CD28 shows a higher peak and faster decline compared to 4-1BB.

Casper, K.M. et al. Nat Rev Clin Oncol 2023; 20: 359-371. Figure recreated using biorender. Maus, M.V. et al. J Immunother Cancer 2020; 8(2):e00111.

Changchong Medically Integrated Oncology: CELEBRATING A DECADE OF IMPACT

23

Baseline Risk Factors for CAR-associated Toxicity

Toxicity Spectrum	Patient	Disease	Product/Tx Related	Additional Considerations
CRS*	Comorbidities +/- age, infections, elevated inflammatory markers, thrombocytopenia	Tumor burden, CNS involvement	Higher dose of CAR-T cells, Co-stimulatory Domain [CD28 > 41BB CARs, Flu-containing/ intensity of LD	Onset/Duration variable and product specific.
ICANS*	CRS, elevated inflammatory markers, younger age, pre-existing neuro conditions, thrombocytopenia	Tumor burden, +/- CNS involvement		ICANS often preceded by CRS
ICAH + hypogam.**	Elevated inflammatory markers, older age, high baseline BM blasts	Refractory, poorly controlled disease, Tumor burden	Multiple prior LOT, intensity of LD	Multifactorial: poor patient condition, tumor burden, pre-treatment LD *Consider CAR-HEMATOTOX
Infections**	Age/comorbidities, Hx of CRS, Existing cytopenias, prior frequent infections, hypogam	Refractory, poorly controlled disease	High-dose & long-duration steroids, other immunosuppressive treatments	Prior infections, biological age vs. chronological age

*Modified EASIX score; LDH>CRP/Platelets is associated with severe CRS and ICANS

**CAR-HEMATOTOX score. Score ≥5 pts are at risk for severe prolonged neutropenia and infections

BM: bone marrow, Hx: history, LOT: lines of therapy, Hypogam: hypogammaglobulinemia, CNS: central nervous system, CRP: c-reactive protein, LDH: lactate dehydrogenase

Reis, A.W. et al. Cancer Medicine 2023; 12: 17. Peng, Y. et al. Biol Blood Marrow Transplant 2019; 25:1029-1037. Garcia, R. et al. Hematology 2019; 18(12): 1205-1210. Saito, M. et al. J Immunother Cancer 2022; 10(12):e0430111. Maus, M.V. et al. J Immunother Cancer 2020; 8(2):e00111.

Changchong Medically Integrated Oncology: CELEBRATING A DECADE OF IMPACT

24

NCODA: Fall Summit

8

Trustworthy Terrain: Navigating Product Choices with Predictable Safety

Best product selection for OP administration is dependent on main 2 factors:

- Attributes of toxicity onset and incidence of severe CRS/ICANS events
 - Delayed onset of toxicities is ideal
 - Potential for OP administration from LD to onset of toxicity warranting IP stay
 - Hospitalization only for toxicity management
 - Low incidences of severe Gr. 2+ CRS/ICANS
 - Management of Gr. 1 CRS in OP setting feasible
- Products with rapid onset of toxicities may be feasible for early discharge options

Example OP CAR-T RWE Studies

Disease State	Product	OP Infusion	Time to Admit (days)	Admit LOS Median (days)
MM, ALL, LBCL	Ida-cel, Tisa-cel, Liso-cel	80% (n=32/40)	2 (pre-emptive D0 approach, n=21)	8-14 vs. 19 IP only
LBCL	Tisa-cel	59% (n=93/157)	2.5 (45% OP required admit)	5 vs. 13 IP only
ALL, LBCL	Axi-cel, Tisa-cel, Brexu-cel	96% (n=47/52)	2-4 (82% OP required admit)	7-10
MM	Cilta-cel	62.8% (n=27/43)	NR (92.6% OP required admit)	4 vs. 11 IP only

Gr: grade, LOS: length of stay RWE: real world experiences, NR: not reported

Chenault CD, et al. Transplant Cellular Ther. 2024; Feb 20(2):131-142. Hansen DM, et al. Front Immunol. 2024; Feb 21; 15:1403532. Kottmann M, et al. Transplant Cell Ther. 2022; Sep 20(2):105-109. Kottmann M, et al. Transplant Cell Ther. 2023; 40(2):77-84. Li, A, et al. Br J Haematol. 2023;233(3):558-562. Winkler, David Hanna, et al. Transplant Cell Ther. 2024; 40(2):1009.

Championing Medically Integrated Oncology: CELEBRATING A DECADE OF IMPACT

25

CRS and ICANS: Incidence, Onset and Duration of FDA Approved CAR T-Cell Products

LBCL Products

	Axi-cel DLBCL ZUMA-1	Axi-cel FL ZUMA-5	Brexu-cel MCL ZUMA-2	Tisa-cel DLBCL JALIS-1	Liso-cel DLBCL TRANSCEND	Liso-cel FL TRANSCEND FL
Gr. 3+ CRS	13%	8%	15%	22%	2%	1%
Median onset/ duration CRS	2 days/ 8 days	4 days/ 8 days	2 days/ 11 days	3 days/ 7 days	5 days/ 5 days	6 days/ 3 days
Gr. 3+ ICANS	28%	21%	31%	12%	10%	2%
Median onset/ duration ICANS	5 days/ 17 days	6 days/ 16 days	7 days/ 12 days	6 days/ 14 days	9 days/ 11 days	8.5 days/ 3.5 days

ALL Products

	Brexu-cel B-ALL ZUMA-3	Tisa-cel B-ALL ELIANA	Cilta-cel B-ALL PULP
Gr. 3+ CRS	24%	48%	3%
Median onset/ duration CRS	5 days/ 7.5 days	3 days/ 8 days	8 days/ 5 days
Gr. 3+ ICANS	25%	13%	7%
Median onset/ duration ICANS	9 days/ 7 days	<8 weeks/ 10 days	12 days/ 8 days

MM Products

	Ida-cel WM Spartan	Cilta-cel WM CARSTITUDE-1
Gr. 3+ CRS	5%	4%
Median onset/ duration CRS	1 day/ 5 days	7 days/ 4 days
Gr. 3+ ICANS	3%	2%
Median onset/ duration ICANS	2 days/ 3 days	8 days/ 4 days

Delayed Onset (~ Day 4+) or low Gr. 3+ events: Potential for OP administration and IP for toxicity management

Early Onset (~ Days 1-3): Potential for early discharge from IP post-toxicity

Hewings DS, et al. N Engl J Med. 2017;377(26):2551-2564. Winkler DS, et al. N Engl J Med. 2019;381(4):435-445. Shaw BD, et al. Lancet. 2021;398(10325). Foster SM, et al. N Engl J Med. 2024; 430(19):2219-2230. Jacobson CA, et al. Lancet Oncol. 2022;23(1). Foster SM, et al. N Engl J Med. 2022;387(26). Kottmann M, et al. Lancet. 2020;395(10249):1015-1024. Kottmann M, et al. Lancet. 2020;395(10249):1015-1024. Kottmann M, et al. N Engl J Med. 2023;389(15):1331-1342. Hewings DS, et al. Lancet. 2021;397(10275). Robinson-Dunn SP, et al. N Engl J Med. 2023;389(11):1020-1034. Sen-Parguel J, et al. N Engl J Med. 2023;389(1), and S, et al. Blood. 2024;144(suppl 1):302.

Championing Medically Integrated Oncology: CELEBRATING A DECADE OF IMPACT

26

Institutional Infrastructure Needs for OP Administration: Dedicated Space or Bed Availability

1

2

3

4

5

6

Clearance for LD

LD Chemo (~3 days)

Administration in OP setting

CAR-T Infusion

Discharge to clinic

(Return Home) ~14-30 days post-infusion

IP stay through toxicity window

Admission to IP Setting for Toxicity Management

Manage low grade toxicities OP

Continuous Daily Visits in OP Setting

Administration in IP setting from LD/or after cell infusion (D0)

Pre-emptive monitoring

Championing Medically Integrated Oncology: CELEBRATING A DECADE OF IMPACT

27

Toxicity Strategies for Outpatient CAR-T Administration




2019 NCODA INTERNATIONAL FALL SUMMIT

28

Considerations for CAR-T Toxicity Monitoring in OP Setting

- Utilize consult services at baseline to identify existing concerns
 - Infectious disease, social work, neurology, etc.
- Create standardized EMR form for daily assessment (CRS/ICANS)
 - Documentation is vital for effective and timely interventions
- Develop institutional criteria/monitoring protocols that are product/disease specific
 - Eligibility for OP CAR-T vs. early transition to OP
 - Create transition of care email group with all stakeholders
- Standardize monitoring requirements during OP CAR-T visits and at home



Olson DD, et al. Transplant Cellular Ther. 2024. Feb 20(2):131-142. Championing Medically Integrated Oncology: CELEBRATING A DECADE OF IMPACT

29

Protocol Considerations for Prompt Escalation of Care

Determine institutional guidance of expected AE's that can be managed in OP setting

- Low-grade CRS, cytopenia's, neutropenic fever, etc.

Develop required IP treatment/monitoring Criteria

- Any Gr. 2 or higher CRS or any ICANS should trigger admission (obs status)



Streamline intake process for IP admission when needed

- Transparency of clinical status: EMR handoff tools, sign-outs, etc.
- Electronic notification tools including all stakeholders
 - Chat/email groups for timely notification

Olson DD, et al. Transplant Cellular Ther. 2024. Feb 20(2):131-142. Championing Medically Integrated Oncology: CELEBRATING A DECADE OF IMPACT

30



ASTCT Consensus CRS Grading and Manifestations

Parameter	Grade 1	Grade 2	Grade 3	Grade 4
Fever*	Temp ≥ 38 °C (100.4 °F)			
	With either:			
Hypotension	None	Not requiring vasopressors	One vasopressor +/- vasopressin	Multiple vasopressors (excluding vasopressin)
	And/or			
Hypoxia	None	Low-flow nasal cannula or blow-by	High-flow nasal cannula, facemask, nonrebreather mask, or Venturi mask	Positive pressure (CPAP, BiPAP, intubation, and mechanical ventilation)

*Fever defined as temperature 38 °C not attributable to any other cause.
• CRS patients who receive antipyretic or anti-cytokine therapy (tocilizumab, steroids, others) **fever is no longer required** to grade subsequent CRS severity
• CRS grading driven by hypotension and/or hypoxia.

Other: Hepatic, renal, cardiovascular, pulmonary, GI, musculoskeletal, neurologic, hematologic
Note: Organ toxicities associated with CRS can be graded per CTCAE v5 but doesn't influence CRS grading

Can progress to life-threatening vasodilatory shock, capillary leak, hypoxia and multiorgan failure

Low-flow nasal cannula: oxygen delivered at ≤ 6 L/minute
High-flow nasal cannula: oxygen delivered at >6 L/minute

Lee DW et al. *Biol Blood Marrow Transplant*. 2019;25(1):625-38

Championing Medically Integrated Oncology: CELEBRATING A DECADE OF IMPACT

31



ASTCT Consensus ICANS Grading and Manifestations

ICE Score		Grade 1	Grade 2	Grade 3	Grade 4
Task	Points	7-9	3-6	0-2	0 (patient is unarousable and unable to perform ICE)
Orientation: orientation to year, month, city, hospital	4	Awakens spontaneously	Awakens to voice	Awakens tactile stimulus only	Unarousable or requires vigorous or repetitive stimuli to arouse. Shunt or coma
Naming: ability to name 3 objects	3	N/A		Any clinical seizure or nonconvulsive seizures on EEG that resolve with intervention	Life-threatening prolonged seizure (>5 min) or repetitive clinical or electrical seizures without return to baseline
Following commands: ability to follow simple commands (e.g., show me the TV remote)	1	N/A			
Writing: ability to write a standard sentence	1	Deep focal motor weakness (hemiparesis or paraparesis)			
Attention: ability to count backwards from 100 by 10	1	N/A		Focal/focal edema on neuroimaging	Diffuse cerebral edema on neuroimaging; or cranial nerve palsy; or papilledema; or Cushing's triad

*A patient with an ICE score of 0 may be classified as grade 3 ICANS if awake with global aphasia, but patient with an ICE of 0 may be classified as grade 4 ICANS if unarousable
%Depressed level of consciousness should be attributable to no other cause (e.g., no sedating medication)

ICE score: Immune Effector Cell-Associated Encephalopathy
ICP: Intracranial pressure, ICE: Immune effector cell associated Encephalopathy

Lee DW et al. *Biol Blood Marrow Transplant*. 2019;25(1):625-38

Championing Medically Integrated Oncology: CELEBRATING A DECADE OF IMPACT

32



Principles of Toxicity Management: No Universal Guideline for Toxicity Management

Protocols vary by institution

CRS: Immune activation correlates with CAR expansion and elevations inflammatory markers and cytokines

ICANS

- Rates of CRS/ICANS vary among products, disease, and patient characteristics
- Appropriate screening per institutional standards
- Baseline labs: CRP, ferritin, CBC/CMP, coagulopathy, and TLS labs

- Symptomatic management of mild CRS is universal and relies on supportive care measures
- **Tocilizumab: 1st line treatment for CRS**
- Steroids typically reserved for tocilizumab-refractory CRS
- Alternative anti-cytokine therapy may be needed in refractory setting

- Tocilizumab not effective
 - Does not penetrate CSF
 - May increase IL-6 levels and worsen ICANS
- **Corticosteroids: 1st line treatment**

CRS: complete blood count, CMP: comprehensive metabolic panel, CSF: cerebrospinal fluid, IL: Interleukin

Sentman CL et al. *Cancer Discov*. 2019 Aug;9(8):959-971. Hay KA, et al. *Blood*. 2017;130(21):2295-2308. Maus MV, et al. *J Immunother Cancer*. 2020;8(2):e001911.

Championing Medically Integrated Oncology: CELEBRATING A DECADE OF IMPACT

33

Toxicity Algorithms and Interventions

CRS onset usually within 1st week after CAR T-cell therapy (peaks within 1-2 weeks)

- #1 is to rule out other etiologies: infections vs. comorbidity flare vs. drug induced

ICANS can be biphasic and occur during CRS and/or present as late or delayed events

- Patients with ICANS should receive AED (if not given prophylactically) + appropriate CNS imaging/EEG monitoring

Gr.	CRS	ICANS	CRS + ICANS
1	Scheduled APAP x 24hrs ± Toci*	Supportive care (± dex)*	Supportive care (± Toci)*
2	Toci ± Dex	Steroids (dex)	Toci + dex
3	Toci + steroid* ± ICU transfer†	Steroids (dex or MP); Consider ICU transfer	Toci + dex or MP
4	Toci + steroid‡§ ± ICU transfer	High-dose MP until Gr 1; ICU/critical care=	Toci + High-dose MP, ICU/critical care=

Steroid dosing: dexamethasone (10-20mg), HD steroids: MP (500-1000mg)

Special Considerations:

- Toci for febrile and those with persistent fever > 39hrs
- † Toci (IV; dex) for high-risk for progression to severe toxicities (i.e. Burden, CD28 prod, age/comorbidities)
- ‡ Consider refractory (MP IV with rapid taper) after 1 or 2 anti-cytokine therapy (e.g. anakinra or etanercept)
- § If life threatening, consider use of anakinra, etanercept, anti-interleukin-6 or IV chemo

Lee DW et al. Blood 2014;124:1188-92, Lee DW et al. Blood 2016;127:4538-4539, Neelapu S et al. J Clin Oncol 2018;36:300-307, Zhou X et al. Blood 2018;131:143-147, Borchmann P et al. J Clin Oncol 2017;35:1547-1553, Zhou X et al. Blood 2018;131:143-147, Borchmann P et al. J Clin Oncol 2017;35:1547-1553, Zhou X et al. Blood 2018;131:143-147, Borchmann P et al. J Clin Oncol 2017;35:1547-1553

AED: antiepileptic drug, EEG: electroencephalogram, toci: ticlopidine, HD: high-dose, MP: methylprednisolone

Championing Medically Integrated Oncology: CELEBRATING A DECADE OF IMPACT

[illegible][illegible]

Risk Adaptive Approaches To Mitigating Severe Toxicities

Risk-Adapted dosing: dosing based on disease burden and/or splitting cell doses

Obe-cel (CAT19-CD19) CAR-T

Holding of infusion if CRS/ICANS

- Real-time dose modifications

Adjusting affinity of binding domain for controlled antigen stimulation

Obe-cel CAT19 scFv

Future Directions:

- Controlled T-cell proliferation 4th and 5th generation CARs
- Logic-gating techniques

Prophylactic/Pre-emptive steroids or anti-cytokine therapy

Dexamethasone (corticosteroid)

Tocilizumab (IL-6R antagonist)

Siltuximab (IL-6 antagonist)

Anakinra (IL-1R antagonist)

Emapalumab (INFγ inhibitor)

Infliximab (TNFα inhibitor)

Bur TA, et al. *Cellular Immunology*. 2024;105:105288. Parvizi AV, et al. *Cancer Metastasis Rev*. 2024 Dec;15(4):1-17.



QUESTION ARS # 2

2025 NCODA INTERNATIONAL
FALL SUMMIT

What prophylactic strategies can be used for high-risk CAR-T patients?

- a. Tocilizumab + dexamethasone given Day -2
- b. Dexamethasone given on Day 0, +1 and +2
- c. There are no prophylaxis strategies at this time as it may impact CAR-T efficacy
- d. Ibrutinib given on Day 0, +1 and +2

Championing Medically Integrated Oncology: CELEBRATING A DECADE OF IMPACT

37

Early Intervention or Prophylactic Strategies to Mitigate CRS Toxicities without Compromising Efficacy

Prophylaxis	Pre-emptive Intervention (as early as Gr. 1)	Concurrent TKI in LBCL
• Tocilizumab (Day 0 or Day +2) • Dexamethasone 10 mg (Days 0, +1 and +2) • Anakinra given on (Days 0, +1 and +2) or extend up to Day +7 • Anakinra given at 1st fever (or Day +2), up to minimum of 10 days	• Dexamethasone • Tocilizumab • Anakinra • Combination of Toci + Dex	- Ibrutinib (BTK Inhibitor) - Icatinib (JAK 1 inhibitor) - Pirtrotinib (BTK Inhibitor)**
		**NCT05658572 in MCL treated with Ibrux+cel

**NCT06553872 in MCL treated with brexu-cel

Chaves OC, et al. *Br Haematol*. 2023;194:650-660. PMID: [37011162](#), [37011163](#), [37011164](#), [37011165](#), [37011166](#), [37011167](#), [37011168](#), [37011169](#), [37011170](#), [37011171](#), [37011172](#), [37011173](#), [37011174](#), [37011175](#), [37011176](#), [37011177](#), [37011178](#), [37011179](#), [37011180](#), [37011181](#), [37011182](#), [37011183](#), [37011184](#), [37011185](#), [37011186](#), [37011187](#), [37011188](#), [37011189](#), [37011190](#), [37011191](#), [37011192](#), [37011193](#), [37011194](#), [37011195](#), [37011196](#), [37011197](#), [37011198](#), [37011199](#), [37011200](#), [37011201](#), [37011202](#), [37011203](#), [37011204](#), [37011205](#), [37011206](#), [37011207](#), [37011208](#), [37011209](#), [37011210](#), [37011211](#), [37011212](#), [37011213](#), [37011214](#), [37011215](#), [37011216](#), [37011217](#), [37011218](#), [37011219](#), [37011220](#), [37011221](#), [37011222](#), [37011223](#), [37011224](#), [37011225](#), [37011226](#), [37011227](#), [37011228](#), [37011229](#), [37011230](#), [37011231](#), [37011232](#), [37011233](#), [37011234](#), [37011235](#), [37011236](#), [37011237](#), [37011238](#), [37011239](#), [37011240](#), [37011241](#), [37011242](#), [37011243](#), [37011244](#), [37011245](#), [37011246](#), [37011247](#), [37011248](#), [37011249](#), [37011250](#), [37011251](#), [37011252](#), [37011253](#), [37011254](#), [37011255](#), [37011256](#), [37011257](#), [37011258](#), [37011259](#), [37011260](#), [37011261](#), [37011262](#), [37011263](#), [37011264](#), [37011265](#), [37011266](#), [37011267](#), [37011268](#), [37011269](#), [37011270](#), [37011271](#), [37011272](#), [37011273](#), [37011274](#), [37011275](#), [37011276](#), [37011277](#), [37011278](#), [37011279](#), [37011280](#), [37011281](#), [37011282](#), [37011283](#), [37011284](#), [37011285](#), [37011286](#), [37011287](#), [37011288](#), [37011289](#), [37011290](#), [37011291](#), [37011292](#), [37011293](#), [37011294](#), [37011295](#), [37011296](#), [37011297](#), [37011298](#), [37011299](#), [37011300](#), [37011301](#), [37011302](#), [37011303](#), [37011304](#), [37011305](#), [37011306](#), [37011307](#), [37011308](#), [37011309](#), [37011310](#), [37011311](#), [37011312](#), [37011313](#), [37011314](#), [37011315](#), [37011316](#), [37011317](#), [37011318](#), [37011319](#), [37011320](#), [37011321](#), [37011322](#), [37011323](#), [37011324](#), [37011325](#), [37011326](#), [37011327](#), [37011328](#), [37011329](#), [37011330](#), [37011331](#), [37011332](#), [37011333](#), [37011334](#), [37011335](#), [37011336](#), [37011337](#), [37011338](#), [37011339](#), [37011340](#), [37011341](#), [37011342](#), [37011343](#), [37011344](#), [37011345](#), [37011346](#), [37011347](#), [37011348](#), [37011349](#), [37011350](#), [37011351](#), [37011352](#), [37011353](#), [37011354](#), [37011355](#), [37011356](#), [37011357](#), [37011358](#), [37011359](#), [37011360](#), [37011361](#), [37011362](#), [37011363](#), [37011364](#), [37011365](#), [37011366](#), [37011367](#), [37011368](#), [37011369](#), [37011370](#), [37011371](#), [37011372](#), [37011373](#), [37011374](#), [37011375](#), [37011376](#), [37011377](#), [37011378](#), [37011379](#), [37011380](#), [37011381](#), [37011382](#), [37011383](#), [37011384](#), [37011385](#), [37011386](#), [37011387](#), [37011388](#), [37011389](#), [37011390](#), [37011391](#), [37011392](#), [37011393](#), [37011394](#), [37011395](#), [37011396](#), [37011397](#), [37011398](#), [37011399](#), [37011400](#), [37011401](#), [37011402](#), [37011403](#), [37011404](#), [37011405](#), [37011406](#), [37011407](#), [37011408](#), [37011409](#), [37011410](#), [37011411](#), [37011412](#), [37011413](#), [37011414](#), [37011415](#), [37011416](#), [37011417](#), [37011418](#), [37011419](#), [37011420](#), [37011421](#), [37011422](#), [37011423](#), [37011424](#), [37011425](#), [37011426](#), [37011427](#), [37011428](#), [37011429](#), [37011430](#), [37011431](#), [370](#)

38

Role of **CAR-HAEMATOTOX (HT)** score and Opportunity for Early Intervention/Supportive Care

Clinical utility of CAR-HT as a validated predictive tool

- Baseline patient characteristics used to evaluate for risk of CRS/ICANS + other complications
 - Risk stratification for severe infections, neutropenia and disease progression
 - The ALL HT omits ferritin for disease burden
- Online calculator for CAR-HT: <https://www.european-mcl.net/home/scores-car-hematotox-286.html>

Anticipate patient needs/allocating resource and guiding toxicity management

- Potential role in guiding escalation of anti-infective prophylaxis (+/- use early G-CSF use and stem cell boost)
- Role for prophylactic strategies (day 0-2) in high-risk patients
- Plan for early intervention strategies

CAR-HT Scoring			
Baseline features	0 points	1 point	2 points
Platelet count	>175K	75-175K	<75K
ANC	≥1200	<1200	-
Hemoglobin	≥9.0	<9.0	-
CRP	<3.0	≥3.0	-
Ferritin	<650	650-2000	>2000
Low risk (HT ^{low}): 0-1 high risk		(HT ^{high}): ≥2 (maximum 7)	

ALL-HT Scoring			
Baseline features	0 points	1 point	2 points
Platelet count	> 175k	75-175k	< 75k
ANC	> 1200	< 1200	
Hgb	> 9	< 9	
CRP	< 3	> 3	
BM blasts	< 5%	5-25%	> 25%
Low risk: 0-3 points		High risk: ≥ 4 points	

k: thousand, G-CSF: Granulocyte Colony-Stimulating Factor

Championing Medically Integrated Oncology: CELEBRATING A DECADE OF IMPACT

39



General Monitoring Strategies for Acute Toxicity

Monitoring for CRS: onset usually within 1st week

- Close hemodynamic monitoring
- Monitor labs for organ dysfunction and inflammatory markers (ferritin/CRP)
- Initiate a full infectious workup and rapid implementation of anti-infective agents upon first signs of fever

Monitoring for ICANS: Manifestations can vary by product and occur up to Day +30

- Waning and waning of symptoms
- ICE assessments at least twice daily (~14 days)

Tools to increase touchpoints

- Telehealth visits
- Post-visit phone calls
- Remote patient monitoring devices
 - Capacity to continually monitor temperature, pulse, respiratory rate and O₂ saturation
 - Can be utilized to decrease caregiver burden

Przy N, Porter D. Biol Blood Marrow Transplant. 2019;25(12):e137-e137. Epub 2019; 142 (Supplement 1).

2007, Lee H. Biol Blood Marrow Transplant. 2019;25(12):e137. Epub 2019; 142 (Supplement 1).

2007, Lee H. Biol Blood Marrow Transplant. 2019;25(12):e137. Epub 2019; 142 (Supplement 1).

2007, Lee H. Biol Blood Marrow Transplant. 2019;25(12):e137. Epub 2019; 142 (Supplement 1).

SOC: standard of care, O₂: oxygen

Championing Medically Integrated Oncology: CELEBRATING A DECADE OF IMPACT

40



General Monitoring/Supportive Care Strategies for Delayed Toxicity

- Late onset toxicities are typically managed by treating facility or locally
 - Supportive care strategies can be instrumental to reduce morbidity/mortality
- B-cell Aplasia/Hypogammaglobulinemia
 - Check IgG levels monthly, then Q3-6mo
 - Administer IVIG if IgG levels < 400 mg/dL or < 600 mg/dL + recurrent infections
- Hematologic toxicity with prolonged cytopenia's (ICAH^T)
 - Etiology multifactorial; Categorized as mild (Gr. 1), moderate (Gr. 2), severe (Gr. 3) and life-threatening (Gr. 4) categories
 - Managed with growth factor + other therapies
- Late infections
 - Ensure empiric antimicrobial are prescribed during neutropenia
 - As early as clearance day

B-cell aplasia/hypogam.

Prolonged cytopenia's

Late infections

Long-term non-ICAH^T toxicity*

Przy N, et al. Biol. Blood Marrow Transplant. 2023;29(12):e137-e137. Epub 2023; 142 (Supplement 1).

2007, Lee H. Biol Blood Marrow Transplant. 2019;25(12):e137. Epub 2019; 142 (Supplement 1).

2007, Lee H. Biol Blood Marrow Transplant. 2019;25(12):e137. Epub 2019; 142 (Supplement 1).

2007, Lee H. Biol Blood Marrow Transplant. 2019;25(12):e137. Epub 2019; 142 (Supplement 1).

ICAH^T: Immune Effector Cell-Associated Hematotoxicity
*Product specific risk most associated with BCMA-directed CAR-T

Championing Medically Integrated Oncology: CELEBRATING A DECADE OF IMPACT

41

Supportive Care Medications: Infection and Other Prophylaxis

TLS/Seizure	Antibacterial	Antiviral	PJP prophy recommended	Antifungal
Allopurinol in high-risk patients • Higher BM burden, WBC and/or LDH (≥ 2000 U/L), etc. Seizure prophy Designated day pre-cells (e.g., D -1) • At time of neurologic toxicity	Recommended for patients with high risk of infections • Prophylaxis may be considered once ANC is <500/μL	VZV prophy in all patients • LD up to 6+ months post-CAR-T and/or until CD4+ count is >0.2 × 10⁹/L	PJP prophy recommended • LD or D+30 up to 6+ months post-CAR-T and/or until CD4+ count is >0.2 × 10⁹/L	Antifungal prophy recommended • Fluconazole/micafungin once ANC is <500/μL • Mold-active azole; high risk of IFI • High CAR-HT score, history of IFI, prolonged neutropenia, use of anti-cytokine therapy +/- steroids

Jan T, et al. Biol. Blood Marrow Transplant. 2023;29(12):e137-e137. Epub 2023; 142 (Supplement 1).

2007, Lee H. Biol Blood Marrow Transplant. 2019;25(12):e137. Epub 2019; 142 (Supplement 1).

2007, Lee H. Biol Blood Marrow Transplant. 2019;25(12):e137. Epub 2019; 142 (Supplement 1).

2007, Lee H. Biol Blood Marrow Transplant. 2019;25(12):e137. Epub 2019; 142 (Supplement 1).

Prophylaxis: prophylaxis, Infection: infection, VZV: varicella virus, PJP: pneumocystis jirovecii pneumonia, IFI: invasive fungal infection

Championing Medically Integrated Oncology: CELEBRATING A DECADE OF IMPACT

42

NCODA: Fall Summit

14


Summary of Strategies to Improve Toxicities in OP Setting

**Risk-adapted Toxicity Management Protocols:** Development of SOP's for early detection and intervention of CRS/ICANS

**Patient/Product Specific Approach:** Assess individualized risk of severe complications + predictability of toxicities to promote delivery of CAR-T in OP setting while preserving patient-centric care

**Optimize Resource Utilization:** Minimize cost burden by favoring OP CAR-T delivery and accelerated discharge pathways while limiting IP stays to manage severe toxicities where possible

43




44


Key Roles of OP Multidisciplinary Team

Specialty	Key Roles/Responsibilities in OP Setting
Nursing	• Provide day to day clinical monitoring and supportive care services • Recognize early CAR-T associated toxicities • Effectively communicate all aspects of CAR T-cell therapy to patients and families, and answer patient questions
APPs/ Providers	• Recognize the unique needs of patients receiving CAR T-cell therapy • Follow CRS/ICANS management algorithms, and administer interventions early • Communicate timely and efficiently with all CAR-T team members and consultative services
Case Managers	• Ensure support for high-cost medications • Assist with transitions of care from OP to IP when needed • Arrange supportive services including home health, PT/OT, etc.
Social workers	• Arrange lodging, transportation, and reimbursement assistance • Provide emotional support to patients and caregivers


45



46



47



48

QUESTION ARS #3

2025 NCODA INTERNATIONAL FALL SUMMIT

Which of the following components is MOST critical to ensuring patient safety in an outpatient CAR-T program?

a. Clearly defined roles for each member of the multidisciplinary team

b. Standardized workflow for patient monitoring and escalation

c. Established protocols for managing cytokine release syndrome and neurotoxicity

d. All of the above

Championing Medically Integrated Oncology: CELEBRATING A DECADE OF IMPACT

49



Best Practices for Institutional Operational Considerations

Regular Meetings

- Include clinical and ancillary members
- Cost/benefit review
- Workflow re-eval

Education of Support Service

- Emergency Department
- Urgent Care
- ICU

Development of SOP's and Workflows

- Patient selection
- Monitoring frequency
- Admission procedure
- Toxicity Management

Staff Education

- Clinical Standards
- Grand rounds
- Lecture series
- Onboarding boot camp for new hires

Quality Control/Metrics

- EMR/SOP variances
- Safety/efficacy Analysis
- Drug-use evaluations
- CIBMTR Reporting
- FACT compliance

Eval: evaluation, ICU: intensive care unit, FACT: Foundation for the Accreditation of Cellular Therapy

Alonso M, et al. Future Oncol. 2023;21(10):1137-1144. Alexander M, et al. Transplant Cell Ther. 2021;27:558-570. Gelleraud A, et al. Clin Hematol Int. 2024;8(2):1-10.

Championing Medically Integrated Oncology: CELEBRATING A DECADE OF IMPACT

50

Multidisciplinary Standard Operating Procedures and Guideline Development



Devise comprehensive procedures + product-specific guidelines for OP

Leverage shared digital platforms for easy access to latest protocol



Complete detailed education for all staff on recognition of CAR-T toxicities and AE management

Review procedures of care escalation

Create process to recognize and intervene with prophylactic measures

Provide refresher training (new hires and cross covering staff)



Attend multidisciplinary meetings and educational sessions to assist with knowledge gaps and skills

Tumor boards

SOP development

Lecture series/Conferences

Participate in QI initiatives



Continually optimize protocols in response to quarterly (or event-driven) protocol-review meetings with key stakeholders

Establish a formal document-control system

Heaven DMO, et al. Front Immunol. 2024, Feb 21;15:1303432. Patel A, et al. Front Immunol. 2024;15:1412002.


QI: quality improvement




51

NCODA: Fall Summit


17



52



53



54



Institution Infrastructure: Triaging and Admissions

Given approximately 30-50% of all OP treated patients may need an admission for CAR-associated toxicities and/or infection complications

- **Ensure adequate staffing for continuity of care**
 - ✓ Dedicated triage line to report after-hours medical issues
 - ✓ Dedicated/knowledge providers on call 24/7
 - ✓ Fellows, residents, hospitalists, APP's
- **Hospital bed access for escalation of care**
 - ✓ Direct admit from OP
 - ✓ Transfer from ED or Urgent Care

Pre-emptive
(Day 0)

Pre-emptive
(Specified day per product)

Toxicity

ED= emergency department

McLean M, et al. Transplant Cell Ther. 2022;26(5):583-589. | Galsword K, et al. Clin Hematol Int. 2024;8(2): 1-20. |
Championing Medically Integrated Oncology: CELEBRATING A DECADE OF IMPACT

55



Patient and Caregiver Education Enhances Safety and Boosts Adherence to Clinical Protocols

- **Step 1:** Effective communication and understanding of caregiver responsibility for close patient monitoring
 - ✓ Need for 24-hour care and proximity to center for at least 2 weeks after treatment
 - ✓ Have a plan to ensure there are no gaps in caregiver availability
- **Step 2:** Patients/caregivers educated about the common AEs of CAR-T and when to seek care or treatment
 - ✓ Both trained in use of thermometer/blood pressure cuff
 - ✓ Provide checklists, wallet cards, etc. in case of transfer from outside facility

Dedicated CAR-T coordinator for 1:1 session

In-person/virtual classes

Prepared materials outlining responsibilities

1:1 clinical pharmacist consult

Education Program

What to do in the event of an adverse event

When to contact

Whom to contact

Where to present

Taylor L, et al. Clin J Oncol Nurs. 2018;22(2):25-35. | Raine M, et al. Future Oncol. 2023;21(10):1137-1144. |
Hansen 2023, et al. Front Immunol. 2024. | Vol. 21-24:35-50/2023. |
Championing Medically Integrated Oncology: CELEBRATING A DECADE OF IMPACT

56

New Changes to REMS Mandate for CAR-T Products: Reducing Barriers to Access and Regulatory Burden

Pros	Cons	Potential Solutions
Removes requirement to maintain special certification and on-site, immediate access to Toc1	Shifts responsibility for CRS/ICANS management onto institution's standard protocols (variable practices)	• Development of SOP's based upon current best practice GL • Robust toxicity surveillance and rapid-response infrastructure
Streamlines product labeling: safety information through BBW/med guides vs. separate REMS program	Without REMS-driven central oversight, data on rare or late toxicities may decline	• Standardize documentation & feedback to improve/support data capture • Ongoing QI metrics

BBW: black box warning, GL: guidelines

FDA Eliminates Risk Evaluation and Mitigation Strategies (REMS) for Autologous Chimeric Antigen Receptor CAR T-cell Immunotherapies. Accessed August 16, 2025. from <https://www.fda.gov/news-events/press-announcements/fda-eliminates-risk-evaluation-and-mitigation-strategies-autologous-chimeric-antigen-receptor-car-t-cell-immunotherapies>. |
Arai YB, et al. Transplant Cell Ther. 2025;29(4):436-437-438-439



57

New Changes to REMS Mandate for CAR-T Products:
Reducing Barriers to Access and Regulatory Burden

Pros	Cons	Potential Solutions
Reduces administrative burden and \$, potential increase in community referrals and improvement in access	Complacency over time, with less frequent refresher training on CAR-T toxicities and mitigation strategies	Use ongoing educational sessions to highlight recognition of CRS and ICANS grading, symptoms and intervention
Shortens patient monitoring windows in updated labels - Driving restrictions 8 → 2 weeks - Proximity requirement 4 → 2 weeks	Removes a formal safety "check" that previously ensured meeting uniform standards	Implement routine virtual check-ins from day 0–14 if not seen in person

FDA Eliminates Risk Evaluation and Mitigation Strategies (REMS) for Autologous Chimeric Antigen Receptor CAR T Cell Immunotherapies. Issued August 15, 2024. From [https://www.fda.gov/oc/2024/08/15/fda-eliminates-rems-autologous-car-t-cell-immunotherapies](#). And 101, et al. "Transfer Out Plan." 2024. [https://doi.org/10.1007/978-1-0717-1247-9](#)

58

Summary: One Step Closer in De-centralizing Access to CAR-T

- OP CAR-T broadens care access, eases burden, and boosts satisfaction
- Enhanced toxicity detection, risk assessment, and early management makes OP delivery feasible
- Eligibility for OP administration hinges on patient and product safety criteria so choose wisely
- Successful implementation requires several components including knowledgeable personnel and clinical space
- Elimination of the REMS requirement is a shift in easing patient/caregiver burden, reducing delivery costs and enhances access equity



59

QUESTION & ANSWER

2025 NCODA INTERNATIONAL FALL SUMMIT


Reimagining CAR-T:
The Outpatient Evolution

Rebecca Gonzalez,
PharmD, BCOP, FASTCT


Blood and Marrow Transplant/Cellular
Immunotherapy Clinical Specialist

Championing Medically Integrated Oncology: CELEBRATING A DECADE OF IMPACT

60



61



62



63
