

Eflapegrastim

Care Team Contact Information: _____

Pharmacy Contact Information: _____

Diagnosis: _____

- Eflapegrastim is used to stimulate the growth of neutrophils (NOO-troh-fils), a type of white blood cell important in the body’s fight against infection.
- Eflapegrastim is often used for people with cancer receiving anti-cancer medicines that can cause fever and a low blood cell count.
- Eflapegrastim may also be used for other reasons.

Goal of Treatment: _____

Treatment Regimen

Treatment Name	How the Treatment Works	How the Treatment is Given
Eflapegrastim (er-LA-peh-gras-tim): Rolvedon (roll-VEH-don)	Helps your body make more white blood cells.	Injection given under the skin in the stomach area (abdomen), back of your arm, or your upper outer buttocks.

Injection Administration and Schedule: Eflapegrastim is typically given once per chemotherapy cycle.

- If you are receiving eflapegrastim because you are also receiving chemotherapy, your last eflapegrastim injection should be at least 14 days before your chemotherapy starts
- Do not get eflapegrastim within 24 hours after your dose of chemotherapy.

Appointments: Appointments may include regular check-ups with your care team, injection appointments, lab visits, and imaging tests. It's important to keep your appointments whenever you can. If you miss any appointments, call your care provider as soon as possible to reschedule your appointment.

Supportive Care to Prevent and Treat Side Effects

Description	Supportive Care Taken at Home
To help prevent bone pain	_____ _____ _____

Review this section if your care team decides that **eflapegrastim injections** can be **given at home** by you or your caregiver:

General Information

- See the detailed “Instructions for Use” that comes with your prescription for information on how to prepare and inject a dose.
- You and your caregiver should be shown how to prepare and inject eflapegrastim before you use it. Talk with your care team.
- If you miss a dose of eflapegrastim, talk to your care team about when you should give your next dose.

Storage Information

- Store eflapegrastim in the refrigerator between 36°F to 46°F (2°C to 8°C).
- Do not freeze eflapegrastim. Throw away (dispose of) any eflapegrastim that has been frozen.
- Store eflapegrastim in the original carton to protect from light.
- Do not shake eflapegrastim.
- Take the carton out of the refrigerator and place the sealed blister tray on a clean flat surface for at least 30 minutes to allow it to reach room temperature before use.
- Throw away (dispose of) any prefilled syringe of eflapegrastim that has been left at room temperature, 68°F to 77°F (20°C to 25°C), for more than 72 hours.
- Keep the eflapegrastim prefilled syringe out of the reach of children and pets.

Common Side Effects

Side Effect	Important Information
Pain in the Bones, Arms, and Legs	<p>Description: After receiving eflapegrastim, you may experience some bone pain. This happens because the medicine helps your bone marrow produce a lot of new white blood cells, which can cause swelling and soreness. You may feel a dull ache in areas like your hips, thighs, back, or arms. This discomfort usually starts 1 to 2 days after your injection and can last for about 2 to 4 days. In some cases, it may continue for up to a week.</p> <p>Recommendations:</p> <ul style="list-style-type: none"> • Your care team may recommend an anti-histamine (such as loratadine (Claritin)) to prevent and treat bone pain. • For some patients, your care team may recommend a nonsteroidal anti-inflammatory drug (NSAID) (such as naproxen (Aleve)) or an alternative pain medicine. Talk with your care team before using these. • These medicines are typically taken for 5 to 7 days after you receive eflapegrastim. <p>Talk to your care team if you have:</p> <ul style="list-style-type: none"> • Bone pain lasting more than 7 days • Severe bone pain

Select Rare Side Effects

Side Effect	Talk to Your Care Team if You Have Any of These Signs or Symptoms
Spleen Rupture	Your spleen may become enlarged and possibly rupture. A ruptured spleen can cause death. <ul style="list-style-type: none"> Pain in the left upper stomach area Pain in your left shoulder
Lung Problems	<ul style="list-style-type: none"> Cough Shortness of breath Chest pain
Sickle Cell Crisis	If you have a sickle cell disorder and receive eflapegrastim, you may have a serious sickle cell crisis. <ul style="list-style-type: none"> Pain Difficulty breathing
Kidney Injury	<ul style="list-style-type: none"> Swelling of your face or ankles Blood in your urine or dark-colored urine You urinate less than usual
Low Platelet Count	<ul style="list-style-type: none"> Bruising easily Frequent nose bleeds Blood in your urine Bood in your spit after a cough
Capillary Leak Syndrome (CLS)	CLS happens when fluids and proteins leak from your small blood vessels into surrounding tissues. It requires immediate treatment to prevent a drop in blood pressure and other serious complications. <ul style="list-style-type: none"> Swelling or puffiness Urinating less than usual Trouble breathing Swelling of your stomach area (abdomen) and feeling of fullness Dizziness or feeling faint A general feeling of tiredness
Inflammation of the Aorta	Inflammation of the aorta (the large blood vessel which transports blood from the heart to the body) has been reported in people who received pegfilgrastim products. <ul style="list-style-type: none"> Fever Abdominal pain Feeling tired Back pain

<p>Bone Marrow Problems called Myelodysplastic Syndrome (MDS) or Acute Myeloid Leukemia (AML)</p>	<p>If you have breast cancer or lung cancer, when eflapegrastim is used with chemotherapy and radiation therapy, or with radiation therapy alone, you may have an increased risk of developing myelodysplastic syndrome (MDS) or acute myeloid leukemia (AML).</p> <ul style="list-style-type: none"> • Weakness • Weight loss • Fever • Frequent infections • Blood in urine or stool • Shortness of breath • Feeling very tired • Bruising or bleeding more easily
<p>Allergic Reactions</p>	<ul style="list-style-type: none"> • Rash over your whole body • Shortness of breath • Wheezing • Dizziness • Swelling around your mouth or eyes • Fast heart rate • Sweating

**Before starting treatment, ask your care team when to call 9-1-1 or seek emergency help.
If you experience any new, worsening, or uncontrolled side effects, contact your care team immediately.**

Pregnancy and Breastfeeding

- It is not known if eflapegrastim will harm an unborn baby.
- It is not known if eflapegrastim passes into breast milk.

Additional Information

- **Tell your care team about all the medicines you take.**
This includes prescriptions, over-the-counter drugs, vitamins, and herbal products. Before starting any new medicine, supplement, or vaccine, ask your care team first.
- **This Patient Education Sheet may not describe all possible side effects.**
Call your healthcare provider for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088.

Notes

Updated Date: March 2, 2026

Scan the QR code below to access this education sheet.



Important notice: The Association of Cancer Care Centers (ACCC), Hematology/Oncology Pharmacy Association (HOPA), Network for Collaborative Oncology Development & Advancement, Inc. (NCODA), and Oncology Nursing Society (ONS) have collaborated in gathering information for and developing this patient education guide. This guide represents a brief summary of the medication derived from information provided by the drug manufacturer and other resources.

This guide does not cover all existing information related to the possible uses, directions, doses, precautions, warnings, interactions, adverse effects, or risks associated with this medication and should not substitute for the advice of a qualified healthcare professional. Provision of this guide is for informational purposes only and does not constitute or imply endorsement, recommendation, or favoring of this medication by ACCC, HOPA, NCODA, or ONS, who assume no liability for and cannot ensure the accuracy of the information presented. All decisions related to taking this medication should be made with the guidance and under the direction of a qualified healthcare professional.

Permission: Patient Education Sheets are provided as a free educational resource for patients with cancer and their caregivers in need of concise, easy to understand information about cancer therapy. Healthcare providers are permitted to copy and distribute the sheets to patients as well as direct patients to the Patient Education Sheets website. However, commercial reproduction or reuse, as well as rebranding or reposting of any type, are strictly prohibited without permission of the copyright holders. Permission requests, including direct linking from Electronic Health Records, and licensing inquiries should be emailed to patienteducation sheets@ncoda.org.

Copyright © 2026 by Network for Collaborative Oncology Development & Advancement, Inc. All rights reserved.

PES-574