

## Positive Quality Intervention: Ixazomib (Ninlaro®) In the Treatment of Multiple Myeloma

**Description:** Ixazomib is an oral proteasome inhibitor indicated in combination with lenalidomide and dexamethasone for the treatment of multiple myeloma after at least one prior therapy. This PQI highlights the management, safety, and efficacy of ixazomib.

**Background:** Multiple Myeloma (MM) is an incurable disease resulting from malignant plasma cells. MM is rare in younger patients (median age at diagnosis 72 years). Patients with MM often experience “CRAB” symptoms defined as: hypercalcemia, renal dysfunction, anemia, and bone lesions. Patients generally relapse multiple times and are treated with multi-drug regimens, preferably triplets; common drugs include proteasome inhibitors (PI), immunomodulatory drugs (IMiD), monoclonal antibodies, and steroids. Select patients may be eligible for an autologous stem cell transplant depending on risk factors associated with disease.

Ixazomib is an oral PI that may be used in the relapsed/refractory (R/R) setting in combination with lenalidomide/dexamethasone. Off-label considerations include utilizing ixazomib in the front-line setting regardless of transplant eligibility in combination with lenalidomide and dexamethasone or in the r/r setting with pomalidomide/dexamethasone or with dexamethasone alone. Ixazomib provides an all oral treatment option in patients with MM. In a clinical phase 1/2 trial, front-line ixazomib/lenalidomide/dexamethasone was associated with a very good partial response in 58% of patients. In the R/R setting, ixazomib-containing regimens demonstrated an improvement in response rates and/or progression free survival. The most common side effects associated with ixazomib in  $\geq 20\%$  of patients include: diarrhea, constipation, thrombocytopenia, peripheral neuropathy, nausea, peripheral edema, vomiting, and back pain.

**PQI Process:** Upon receipt of prescription for ixazomib:

- Prior to therapy initiation, obtain baseline labs CBC/diff, CMP, and relevant MM labs to assess treatment response
  - Select MM labs: lactate dehydrogenase (LDH), beta-2 microglobulin, protein analyses, bone marrow aspirate, and cytogenetic studies
  - Platelet and neutrophils nadir occurs on days 14-21 of each cycle
  - Labs should be monitored monthly but may be more frequent with the first three cycles
- Ensure patients receiving PI therapies are also receiving antiviral therapy to prevent against herpes zoster reactivation
- Patients on IMiDs (lenalidomide or pomalidomide) should be on aspirin for DVT/PE prophylaxis or therapeutic anticoagulation based on clotting history and risk factors
- Patients receiving ixazomib are likely to receive IMiD therapy. Ensure REMS requirements are met with IMiD therapy for a timely initiation of treatment and to keep cycles on track
- Ixazomib recommended dosing: 4 mg days 1, 8, 15, every 28 days
  - Dose reduce for severe renal impairment (or ESRD requiring dialysis), or moderate and severe hepatic impairment at baseline.
    - Baseline CrCl  $<30$  ml/min or ESRD requiring dialysis: 3 mg PO daily on days 1, 8, 15 every 28 days
    - Baseline total bilirubin  $>1.5x$  ULN: 3 mg PO daily on days 1, 8, 15 every 28 days
  - If toxicity occurs during treatment, ixazomib should be held and dose reduced to the next lower dose if attributed to therapy
  - Ixazomib is available as 4 mg, 3 mg, and 2.3 mg allowing for 25% incremental dose adjustments
- Avoid concomitant administration of ixazomib with strong CYP3A inducers

**Important Notice:** NCODA has developed this Positive Quality Intervention platform. This platform represents a brief summary of medication uses and therapy options derived from information provided by the drug manufacturer and other resources. This platform is intended as an educational aid and does not provide individual medical advice and does not substitute for the advice of a qualified healthcare professional. This platform does not cover all existing information related to the possible uses, directions, doses, precautions, warning, interactions, adverse effects, or risks associated with the medication discussed in the platform and is not intended as a substitute for the advice of a qualified healthcare professional. The materials contained in this platform are for informational purposes only and do not constitute or imply endorsement, recommendation, or favoring of this medication by NCODA, which assumes no liability for and does not ensure the accuracy of the information presented. NCODA does not make any representations with respect to the medications whatsoever, and any and all decisions, with respect to such medications, are at the sole risk of the individual consuming the medication. All decisions related to taking this medication should be made with the guidance and under the direction of a qualified healthcare professional. *Updated 3.21.22*

<b>Alternating Dose Reductions with Lenalidomide</b>			
<b>Reaction</b>	Platelets	Neutrophils	Rash
<b>Parameter</b>	< 30,000/mm <sup>3</sup>	< 500/mm <sup>3</sup>	Grade 2 or 3
<b>First Occurrence</b>	Hold ixazomib and lenalidomide Upon recovery, resume both but <b>reduce lenalidomide</b> at next lower dose and <b>continue ixazomib</b> at most recent dose		
<b>Subsequent Occurrence</b>	Hold ixazomib and lenalidomide Upon recovery, resume both but <b>reduce ixazomib</b> at next lower dose and <b>continue lenalidomide</b> at most recent dose		

- Monitor for peripheral neuropathy
- It is highly recommended that a therapy calendar be utilized for all patients with MM due to complexity of therapies, older age of patients, and the potential lack of a medically-integrated pharmacy (ex. triplet-drug regimens may arrive from different sources) (see supplemental information for example)

**Patient Centered Activities:**

- Provide [Oral Chemotherapy Education \(OCE\)](#) Sheet and therapy calendar to all patients
- Ensure patients understand the dosing schedule and consider providing a calendar
  - Ixazomib is administered on days 1, 8 and 15 every 28 days
  - Steroid component of regimen may be taken prior to ixazomib to help with nausea control but steroids should be taken with food
  - Ixazomib should be taken on an empty stomach (1 hour before or 2 hours after a meal/snack)
- Advise patients that a missed dose should not be taken with 3 days of the next scheduled dose.
  - If vomiting occurs, do not repeat dose
- Patient Assistance: [NCODA Financial Assistance Tool](#)

**Supplemental Information**

<b>Example 28-Day Dosing Calendar (ixazomib, lenalidomide, dexamethasone)</b>									
	<b>Week 1</b>		<b>Week 2</b>		<b>Week 3</b>		<b>Week 4</b>		
	<b>Day 1</b>	<b>Days 2-7</b>	<b>Day 8</b>	<b>Days 9-14</b>	<b>Day 15</b>	<b>Days 16-21</b>	<b>Day 22</b>	<b>Days 23-28</b>	
<b>Ixazomib</b>	✓		✓		✓				
<b>Lenalidomide</b>	Take every day on days 1-21								
<b>Dexamethasone</b>	✓		✓		✓		✓		

**References:**

1. [Ninlaro® \(ixazomib\) \[prescribing information\]](#).
2. SEER Stat Fact Sheets: Myeloma. Available at <http://seer.cancer.gov/statfacts/html/mulmy.html>.

**Important Notice:** NCODA has developed this Positive Quality Intervention platform. This platform represents a brief summary of medication uses and therapy options derived from information provided by the drug manufacturer and other resources. This platform is intended as an educational aid and does not provide individual medical advice and does not substitute for the advice of a qualified healthcare professional. This platform does not cover all existing information related to the possible uses, directions, doses, precautions, warning, interactions, adverse effects, or risks associated with the medication discussed in the platform and is not intended as a substitute for the advice of a qualified healthcare professional. The materials contained in this platform are for informational purposes only and do not constitute or imply endorsement, recommendation, or favoring of this medication by NCODA, which assumes no liability for and does not ensure the accuracy of the information presented. NCODA does not make any representations with respect to the medications whatsoever, and any and all decisions, with respect to such medications, are at the sole risk of the individual consuming the medication. All decisions related to taking this medication should be made with the guidance and under the direction of a qualified healthcare professional. Updated 3.21.22